


(19)



(11)

EP 1 296 143 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
03.06.2009 Bulletin 2009/23

(51) Int Cl.:
G01N 33/68 (2006.01) G01N 33/58 (2006.01)

(21) Application number: **02025544.4**

(22) Date of filing: **02.09.1998**

(54) Mass spectrometric detection of polypeptides

Massenspektrometrischer Nachweis von Polypeptiden

Détection de polypeptides par spectroscopie de masse

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

(30) Priority: **02.09.1997 US 922201**

(43) Date of publication of application:
26.03.2003 Bulletin 2003/13

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
98943528.4 / 1 010 008

(73) Proprietor: **SEQUENOM, INC.**
San Diego,
California 92121 (US)

(72) Inventors:
• **Higgins, Scott**
Bearsden,
Glasgow GL1 3BB (GB)
• **Köster, Hubert**
6900 Lugano-Cassarate (CH)
• **Little, Daniel P.**
Boston, MA 02110 (US)
• **Lough, David**
Eyemouth, Berwickshire TD1Y 55A (GB)

(74) Representative: **Baldock, Sharon Claire et al**
BOULT WADE TENNANT
Verulam Gardens
70 Gray's Inn Road
London WC1X 8BT (GB)

(56) References cited:
WO-A-93/24834 WO-A-94/28418
WO-A-96/36732 WO-A-96/36986
GB-A- 2 168 478

- **PROME D ET AL: "Use of Combined Mass Spectrometry Methods for the Characterization of a New Variant of Human Hemoglobin: The Double Mutant Hemoglobin Villeparisis beta77 (EF1) His -> Tyr, beta80 (EF4) Asn -> Ser" JOURNAL OF THE AMERICAN SOCIETY FOR MASS SPECTROMETRY, ELSEVIER SCIENCE INC, US, vol. 7, no. 2, 1 February 1996 (1996-02-01), pages 163-167, XP004051911 ISSN: 1044-0305**
- **MOSCA A ET AL: "Hb Abruzzo [beta 143(H21)His -> Arg] Identified by Mass Spectrometry and DNA Analysis" HEMOGLOBIN, XX, XX, vol. 17, no. 3, 1993, pages 261-268, XP002093188**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 296 143 B1

Description**FIELD OF THE INVENTION**

5 **[0001]** The disclosed processes relate generally to the field of proteomics and molecular medicine, and more specifically to processes using mass spectrometry to determine the identity of a target polypeptide.

BACKGROUND

10 **[0002]** In recent years, the molecular biology of a number of human genetic diseases has been elucidated by the application of recombinant DNA technology. More than 3000 diseases are known to be of genetic origin (Cooper and Krawczak, "Human Genome Mutations" (BIOS Publ. 1993)), including, for example, hemophilias, thalassemias, Duchenne muscular dystrophy, Huntington's disease, Alzheimer's disease and cystic fibrosis, as well as various cancers such as breast cancer. In addition to mutated genes that result in genetic disease, certain birth defects are the result of

15 chromosomal abnormalities, including, for example, trisomy 21 (Down's syndrome), trisomy 13 (Patau syndrome), trisomy 18 (Edward's syndrome), monosomy X (Turner's syndrome) and other sex chromosome aneuploidies such as Klinefelter's syndrome (XXY).

[0003] Other genetic diseases are caused by an abnormal number of trinucleotide repeats in a gene. These diseases include Huntington's disease, prostate cancer, spinal cerebellar ataxia 1 (SCA-1), Fragile X syndrome (Kremer et al., Science 252:1711-14 (1991); Fu et al., Cell 67:1047-58 (1991); Hirst et al., J. Med. Genet. 28:824-29 (1991)); myotonic dystrophy type I (Mahadevan et al., Science 255:1253-55 (1992); Brook et al., Cell 68:799-808 (1992)), Kennedy's disease (also termed spinal and bulbar muscular atrophy (La Spada et al., Nature 352:77-79 (1991)), Machado-Joseph disease, and dentatorubral and pallidolysian atrophy. The aberrant number of triplet repeats can be located in any region of a gene, including a coding region, a non-coding region of an exon, an intron, or a regulatory element such as

25 a promoter. In certain of these diseases, for example, prostate cancer, the number of triplet repeats is positively correlated with prognosis of the disease

[0004] Evidence indicates that amplification of a trinucleotide repeat is involved in the molecular pathology in each of the disorders listed above. Although some of these trinucleotide repeats appear to be in non-coding DNA, they clearly are involved with perturbations of genomic regions that ultimately affect gene expression. Perturbations of various

30 dinucleotide and trinucleotide repeats resulting from somatic mutation in tumor cells also can affect gene expression or gene reputation.

[0005] Additional evidence indicates that certain DNA sequences predispose an individual to a number of other diseases, including diabetes, arteriosclerosis, obesity, various autoimmune diseases and cancers such as colorectal, breast, ovarian and lung cancer. Knowledge of the genetic lesion causing or contributing to a genetic disease allows one to

35 predict whether a person has or is at risk of developing the disease or condition and also, at least in some cases, to determine the prognosis of the disease.

[0006] Numerous genes have polymorphic regions. Since individuals have any one of several allelic variants of a polymorphic region, each can be identified based on the type of allelic variants of polymorphic regions of genes. Such identification can be used, for example, for forensic purposes. In other situations, it is crucial to know the identity of allelic

40 variants in an individual. For example, allelic differences in certain genes such as the major histocompatibility complex (MHC) genes are involved in graft rejection or graft versus host disease in bone marrow transplantation. Accordingly, it is highly desirable to develop rapid, sensitive, and accurate methods for determining the identity of allelic variants of polymorphic regions of genes or genetic lesions.

[0007] Several methods are used for identifying of allelic variants or genetic lesions. For example, the identity of an allelic variant or the presence of a genetic lesion can be determined by comparing the mobility of an amplified nucleic acid fragment with a known standard by gel electrophoresis, or by hybridization with a probe that is complementary to the sequence to be identified. Identification, however, only can be accomplished if the nucleic acid fragment is labeled with a sensitive reporter function, for example, a radioactive (^{32}P , ^{35}S), fluorescent or chemiluminescent reporter. Radioactive labels can be hazardous and the signals they produce can decay substantially over time. Non-radioactive

50 labels such as fluorescent labels can suffer from a lack of sensitivity and fading of the signal when high intensity lasers are used. Additionally, labeling, electrophoresis and subsequent detection are laborious, time-consuming and error-prone procedures. Electrophoresis is particularly error-prone, since the size or the molecular weight of the nucleic acid cannot be correlated directly to its mobility in the gel matrix because sequence specific effects, secondary structures and interactions with the gel matrix-cause artifacts in its migration through the gel.

55 **[0008]** Mass spectrometry has been used for the sequence analysis of nucleic acids (see, for example, Schram, Mass Spectrometry of Nucleic Acid Components, Biomedical Applications of Mass Spectrometry 34:203-287 (1990); Crain, Mass Spectrom. Rev. 9:505-554 (1990); Murray, J. Mass Spectrom. Rev. 31:1203 (1996); Nordhoff et al., J. Mass Spectrom. 15:67 (1997)). In general, mass spectrometry provides a means of "weighing" individual molecules by ionizing

the molecules *in vacuo* and making them "fly" by volatilization. Under the influence of electric and/or magnetic fields, the ions follow trajectories depending on their individual mass (m) and charge (z). For molecules with low molecular weight, mass spectrometry is part of the routine physical-organic repertoire for analysis and characterization of organic molecules by the determination of the mass of the parent molecular ion. In addition, by arranging collisions of this parent molecular ion with other particles such as argon atoms, the molecular ion is fragmented, forming secondary ions by collisionally activated dissociation (CAD); the fragmentation pattern/pathway very often allows the derivation of detailed structural information. Many applications of mass spectrometric methods are known in the art, particularly in the bio-sciences (see Meth. Enzymol., Vol. 193, "Mass Spectrometry" (McCloskey, ed.; Academic Press, NY 1990; McLaffery et al., Acc. Chem. Res. 27:297-386 (1994); Chait and Kent, Science 257:1885-1894 (1992); Siuzdak, Proc. Natl. Acad. Sci., USA 91:11290-11297 (1994)), including methods for producing and analyzing biopolymer ladders (see, International PCT application No. WO 96/36732; U.S. Patent No. 5,792,664). GB 2,168,478 discloses a method for determining the addition or deletion of amino acids from the terminal ends of peptides. The method comprises digesting a peptide to produce C- or N- terminus fragments and subjecting these to mass spectrometry to analyse the presence of peptides having an unexpected mass. Despite the effort to apply mass spectrometry methods to the analysis of nucleic acid molecules, however, there are limitations, including physical and chemical properties of nucleic acids. Nucleic acids are very polar biopolymers that are difficult to volatilize.

[0009] Accordingly, a need exists for methods to determine the identity of a nucleic acid molecules, particularly genetic lesions in a nucleic acid molecule, using alternative methodologies. Therefore it is an object herein to provide processes and compositions that satisfy this need and provide additional advantages.

SUMMARY OF THE INVENTION

[0010] Processes for determining the identity of a target polypeptide by mass spectrometry are provided as described in in the claims. Identity includes identifying the sequence of the polypeptide.

[0011] As disclosed herein, the polypeptide can be obtained by methods including transcribing a nucleic acid encoding the target polypeptide into RNA and translating the RNA into the target polypeptide. If desired, transcription of the nucleic acid or translation of the RNA, or both, can be performed *in vitro*.

[0012] A process as disclosed herein also can include a step of amplifying a nucleic acid encoding the target polypeptide, for example, by performing the polymerase chain reaction (PCR) using a forward primer and a reverse primer. The forward primer or the reverse primer can contain an RNA polymerase promoter such as an SP6 promoter, T3 promoter, or T7 promoter. In addition, a primer can contain a nucleotide sequence for a transcription start site. A primer also can encode a translation START (ATG) codon. Accordingly, a target polypeptide can be translated from a nucleic acid that is not naturally transcribed or translated *in vivo*, for example, by incorporating a START codon in the nucleic acid to be translated, thereby providing a translation reading frame. Furthermore, a primer can contain a nucleotide sequence, or complement thereof, encoding a second peptide or polypeptide, for example, a tag peptide such as a myc epitope tag, a *Haemophilus influenza* hemagglutinin peptide tag, a polyhistidine sequence, a polylysine sequence or a polyarginine sequence. A process as disclosed herein can be performed *in vivo*, for example, in a host cell such as a bacterial host cell transformed with a nucleic acid encoding a target polypeptide or a eukaryotic host cell such as a mammalian cell transfected with a nucleic acid encoding a target polypeptide.

[0013] A process as disclosed is performed using a mass spectrometric analysis, including for example, matrix assisted laser desorption ionization (MALDI), continuous or pulsed electrospray ionization, ionspray, thermospray, or massive cluster impact mass spectrometry and a detection format such as linear time-of-flight (TOF), reflectron time-of-flight, single quadrupole, multiple quadrupole, single magnetic sector, multiple magnetic sector, Fourier transform ion cyclotron resonance, ion trap, and combinations thereof such as MALDI-TOF spectrometry. An advantage of using a process as provided is that no radioactive label is required. Another advantage is that relatively short polypeptides can be synthesized from a target nucleic acid, thus providing an accurate measurement of molecular weight by mass spectrometry, as compared to analysis of the nucleic acid itself.

[0014] An RNA molecule encoding a target polypeptide can be translated in a cell-free extract, which can be a eukaryotic cell-free extract such as a reticulocyte lysate, a wheat germ extract, or a combination thereof; or a prokaryotic cell-free extract, for example, a bacterial cell extract such as an *E. coli* S30 extract. If desired, translation and transcription of a target nucleic acid can be performed in the same cell-free extract, for example, a reticulocyte lysate or a prokaryotic cell extract.

[0015] Target polypeptides generally are isolated prior to being detected by mass spectrometric analysis. For example, the polypeptides can be isolated from a cell or tissue obtained from a subject such as a human. The target polypeptide can be isolated using a reagent that interacts specifically with the target polypeptide, for example, an antibody that interacts specifically with the target polypeptide, or the target polypeptide can be fused to a tag peptide and isolated using a reagent that interacts specifically with the tag peptide, for example, an antibody specific for the tag peptide. A reagent also can be another molecule that interacts specifically with the tag peptide, for example, metal ions such as

nickel or cobalt ions, which interact specifically with a hexahistidine (His-6) tag peptide.

[0016] Target polypeptides can be immobilized to a solid support, such as a bead or a microchip, which can be a flat surface or a surface with structures made of essentially any material commonly used for fashioning such a device. A microchip is useful, for example, for attaching moieties in an addressable array. Immobilization of a target polypeptide provides a means to isolate the polypeptide, as well as a means to manipulate the isolated target polypeptide prior to mass spectrometry.

[0017] Methods are provided for sequencing an immobilized target polypeptides, including sequencing from the carboxyl terminus or from the amino terminus. Furthermore, methods of determining the identity of each of the target polypeptides in a plurality of target polypeptides by multiplexing are provided.

[0018] In particular embodiments, post translational capture and immobilization of a target polypeptide via a cleavable linker are provided in order to orthogonally sequence a polypeptide. These methods can include: 1) obtaining the target polypeptide; 2) immobilizing the target polypeptide to a solid surface; 3) treating the immobilized target polypeptide with an enzyme or chemical in a time dependent manner to generate a series of deleted fragments; 4) the cleaved polypeptide fragments are conditioned; 5) cleaving the linker and thereby releasing the immobilized fragments; 6) determining the mass of the release fragments; and 7) aligning the masses of each of the polypeptide fragments to determine the amino acid sequence. Variants of these methods in which one or more steps are combined or eliminated are also contemplated.

[0019] In one embodiment, the second step includes immobilizing the amino terminal portion of the polypeptide to a solid support via a photocleavable linker. In a more preferred embodiment, the solid support is activated as described in Figure 2 and allowed to react with the amino group of a target polypeptide.

[0020] In another embodiment, the second step comprises immobilizing the carboxy terminal portion of the polypeptide to a solid support via a photocleavable linker. In a more preferred embodiment, a photocleavable linker is a linker that can be cleaved from the solid support with light. In a more preferred embodiment, the solid support is activated as described in Figure 3 and allowed to react with the carboxy group of a target polypeptide.

[0021] In another embodiment, the second step includes immobilizing either the carboxy or amino termini of group of different polypeptides to a solid support in an array format via a photocleavable linker. In a more preferred embodiment, discrete areas of a silicon surface are activated with the chemistry described in Figure 2 and an array composed of from 2 to 999 positions.

[0022] In another embodiment, the second step includes immobilizing the amino terminal portion of the polypeptide to a solid support via a cleavable linker. In a more preferred embodiment, a cleavable linker is a silyl linker that can be cleaved from the solid support. In a more preferred embodiment, the solid support is activated as described in Figure 2 and allowed to react with the amino group of a target polypeptide.

[0023] In another embodiment, the second step includes immobilizing the carboxy terminal portion of the polypeptide to a solid support via a cleavable linker. In a more preferred embodiment, a cleavable linker is a silyl linker that can be cleaved from the solid support. In a more preferred embodiment, the solid support is activated as described in Figure 3 and allowed to react with the carboxy group of a target polypeptide.

[0024] In another embodiment, the second step includes immobilizing either the carboxy or the amino termini of group of different polypeptides to a solid support in an array format via a cleavable linker. In a more preferred embodiment, discrete areas of a silicon surface are activated with the chemistry described in Figure 2, thereby forming an array, preferably composed of from 2 to 999 positions.

[0025] In another embodiment, the third step includes immobilizing the amino terminal end of the target polypeptide (s) to the solid support and treating with an exopeptidase. In a preferred embodiment, exopeptidase digestion is carried out in a time dependent manner to generate a nested group of immobilized polypeptide fragments of varying lengths. In a more preferred embodiment, exopeptidase is selected from a group of one or more mono-peptidases and polypeptidases including carboxypeptidase Y, carboxypeptidase P, carboxypeptidase A, carboxypeptidase G and carboxypeptidase B.

[0026] In another embodiment, the exopeptidase is selected from a group of one or more mono-peptidases and polypeptidases including aminopeptidases including alanine aminopeptidase, leucine aminopeptidase, pyroglutamate peptidase, dipeptidyl peptidase, microsomal peptidase and other enzymes which progressive digest the amino terminal end of a polypeptidase.

[0027] In another embodiment, the third step comprises a step where exopeptidase digestion is carried out under reaction conditions that remove any secondary or tertiary structure, leaving the terminal residues of the polypeptide inaccessible to exopeptidases. In a preferred embodiment, the reaction conditions expose the terminus of a target polypeptide(s) to temperatures over about 70 ° C and below about 100 ° C. In a more preferred embodiment, the exopeptidase is a thermostable carboxypeptidase or aminopeptidase. In another preferred embodiment, the reaction conditions expose the terminus of a target polypeptide(s) to high ionic strength conditions. In a more preferred embodiment, the exopeptidase is a salt tolerant carboxypeptidase or aminopeptidase.

[0028] In another embodiment, the second step includes conditioning of polypeptide after enzymatic treatment or purification. In a more preferred embodiment, methods of conditioning include methods that prepare the polypeptide or

polypeptide fragments in a manner that generally improves mass spectrometric analysis. In a more preferred embodiment, conditioning may include cation exchange.

[0029] A target polypeptide can be encoded by an allelic variant of a polymorphic region of a gene of a subject, or can be encoded by an allelic variant of a polymorphic region that is located in a chromosomal region that is not in a gene.

5 A process as disclosed herein can include a step of determining whether the allelic variant is identical to an allelic variant of a polymorphic region that is associated with a disease or condition, thereby indicating whether a subject has or is at risk of developing the disease or condition associated with the specific allelic variant of the polymorphic region of the gene. The disease or condition can be associated, for example, with an abnormal number of nucleotide repeats, for example, dinucleotide, trinucleotide, tetranucleotide or pentanucleotide repeats. Since trinucleotide repeats, for example, 10 can be very long, determination of the number of trinucleotide repeats by analyzing the DNA directly would not be straightforward. Since a process for determining the identity of a target polypeptide as disclosed herein is based on the analysis of a polypeptide, particularly a polypeptide encoded essentially by trinucleotide repeats, determination of the number of trinucleotide repeats will be more accurate using the disclosed processes. A disease or condition that can be identified using a disclosed process includes, for example, Huntington's disease, prostate cancer, Fragile X syndrome 15 type A, myotonic dystrophy type I, Kennedy's disease, Machado-Joseph disease, dentatorubral and pallidolysian atrophy, and spino bulbar muscular atrophy; as well as aging, which can be identified by examining the number of nucleotide repeats in telomere nucleic acid from a subject. The disease or condition also can be associated with a gene such as genes encoding BRCA1, BRCA2, APC; a gene encoding dystrophin, β -globin, Factor IX, Factor VIIc, ornithine-d-aminotransferase, hypoxanthine guanine phosphoribosyl transferase, or the cystic fibrosis transmembrane receptor (CFTR); 20 or a proto-oncogene.

[0030] A process as disclosed herein can be used to genotype a subject by determining the identity of one or more allelic variants of one or more polymorphic regions in one or more genes or chromosomes of the subject. For example, the one or more genes can be associated with graft rejection and the process can be used to determine compatibility between a donor and a recipient of a graft. Such genes can be MHC genes, for example. Genotyping a subject using a 25 process as provided herein can be used for forensic or identity testing purposes and the polymorphic regions can be present in mitochondrial genes or can be short tandem repeats.

[0031] A disclosed process also can be used to determine whether a subject carries a pathogenic organism such as a virus, bacterium, fungus or protist. A process for determining the isotype of a pathogenic organism also is provided. Thus, depending on the sequence to be detected, the processes disclosed herein can be used, for example, to diagnose 30 a genetic disease or chromosomal abnormality; a predisposition to or an early indication of a gene influenced disease or condition, for example, obesity, atherosclerosis, diabetes or cancer; or an infection by a pathogenic organism, for example, a virus, bacterium, parasite or fungus; or to provide information relating to identity, heredity or compatibility using, for example, mini-satellite or micro-satellite sequences or HLA phenotyping.

[0032] A process as disclosed herein provides a means for determining the amino acid sequence of a polypeptide of interest. Such a process can be performed by using mass spectrometry to determine the identity of an amino acid residue released from the amino terminus or the carboxyl terminus of a polypeptide of interest. Such a process also can be performed, for example, by producing a nested set of carboxyl terminal or amino terminal deletion fragments of a polypeptide of interest, or peptide fragment thereof, and subjecting the nested set of deletion fragments to mass spec- 35 trometry, thereby determining the amino acid sequence of the polypeptide.

40 **[0033]** A process of determining the amino acid sequence of a polypeptide of interest can be performed, for example, using a polypeptide that is immobilized, reversibly, if desired, to a solid support. In addition, such a process can be performed on a plurality of such polypeptides, which can be, for example, a plurality of target polypeptides immobilized in an addressable array on a solid support such as a microchip, which can contain, for example, at least 2 positions, and as many as 999 positions, or 1096 positions, or 9999 positions, or more. In general, a target polypeptide, or the 45 amino acids released therefrom, are conditioned prior to mass spectrometry, thereby increasing resolution of the mass spectrum. A target polypeptide is conditioned by mass modification. In addition, the amino acid sequences of a plurality of mass modified target polypeptide can be determined by mass spectrometry using a multiplexing format.

BRIEF DESCRIPTION OF THE DRAWINGS

50 **[0034]**

FIGURE 1A shows the nucleotide sequence of a nucleic acid (SEQ ID NO: 8) that can be obtained by PCR ampli- 55 fication of DNA containing a non-variable stretch of 12 CAG repeats (shown without italics) and a variable repeat of 10 CAG repeat units (represented in italics) with primers (underlined) having the sequence (forward primer) or the complement of the sequence (reverse primer). The T7 promoter sequence and the sequence encoding a hexahistidine (His-6) peptide are represented in bold.

FIGURE 1B shows the sequence (SEQ ID NO: 9) of the 71 amino acid polypeptide encoded by the nucleic acid

sequence shown in Figure 1A. The stretch of 10 variable glutamine (Q) residues encoded by the trinucleotide repeats is represented in italics. The His-6 peptide is represented in bold.

FIGURE 2 sets forth an exemplary scheme for orthogonal capture, cleavage and MALDI analysis of a polypeptide. The peptide is conjugated to a solid surface, which can be a microchip, through the use of an acid cleavable diisopropylsilyl linker. The peptide is conjugated to the linker at its amino terminus through the formation of an amide bond. The immobilized polypeptide can be truncated, for example, using a carboxypeptidase, or can be cleaved using an endopeptidase such as trypsin, then is cleaved from the solid support by exposure to acidic conditions such as the 3-HPA (3-hydroxypicolinic acid) matrix solution. The cleaved polypeptide then is subjected to mass spectrometry, for example, MALDI.

FIGURE 3 illustrates additional linkers and capture strategies for reversibly immobilizing a polypeptide on a solid surface. Figure 3 provides reaction conditions for conjugating a polypeptide by its carboxyl terminus to a solid support using 1-ethyl-3-(3-dimethylamino-propyl) carbodiimide hydrochloride (EDC)/N-hydroxy succinimidyl (NHS).

DETAILED DESCRIPTION OF THE INVENTION

DEFINITIONS

[0035] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which this invention belongs. For convenience, the meaning of certain terms and phrases used in the specification and claims are provided.

[0036] As used herein, the term "allele" refers to an alternative form of a nucleotide sequence in a chromosome. Reference to an "allele" includes a nucleotide sequence in a gene or a portion thereof, as well as a nucleotide sequence that is not a gene sequence. Alleles occupy the same locus or position on homologous chromosomes. A subject having two identical alleles of a gene is considered "homozygous" for the allele, whereas a subject having two different alleles is considered "heterozygous." Alleles of a specific nucleotide sequence, for example, of a gene can differ from each other in a single nucleotide, or several nucleotides, where the difference can be due to a substitution, deletion, or insertion of one or more nucleotides. A form of a gene containing a mutation is an example of an allele. In comparison, a wild-type allele is an allele that, when present in two copies in a subject, results in a wild-type phenotype. There can be several different wild-type alleles of a specific gene, since certain nucleotide changes in a gene may not affect the phenotype of a subject having two copies of the gene with the nucleotide changes.

[0037] The term "allelic variant" refers to a portion of an allele containing a polymorphic region in the chromosomal nucleic acid. The term "allelic variant of a polymorphic region of a gene" refers to a region of a gene having one of several nucleotide sequences found in that region of the gene in different individuals. The term "determining the identity of an allelic variant of a polymorphic region" refers to the determination of the nucleotide sequence or encoded amino acid sequence of a polymorphic region, thereby determining to which of the possible allelic variants of a polymorphic region that particular allelic variant corresponds.

[0038] The term "polymorphism" refers to the coexistence, in a population, of more than one form of an allele. A polymorphism can occur in a region of a chromosome not associated with a gene or can occur, for example, as an allelic variant or a portion thereof of a gene. A portion of a gene that exists in at least two different forms, for example, two different nucleotide sequences, is referred to as a "polymorphic region of a gene." A polymorphic region of a gene can be localized to a single nucleotide, the identity of which differs in different alleles, or can be several nucleotides long.

[0039] As used herein, the term "biological sample" refers to any material obtained from a living source, for example, an animal such as a human or other mammal, a plant, a bacterium, a fungus, a protist or a virus. The biological sample can be in any form, including a solid material such as a tissue, cells, a cell pellet, a cell extract, or a biopsy, or a biological fluid such as urine, blood, saliva, amniotic fluid, exudate from a region of infection or inflammation, or a mouth wash containing buccal cells.

[0040] The term "polypeptide," as used herein, means at least two amino acids, or amino acid derivatives, including mass modified amino acids, that are linked by a peptide bond, which can be a modified peptide bond. A polypeptide can be translated from a nucleotide sequence that is at least a portion of a coding sequence, or from a nucleotide sequence that is not naturally translated due, for example, to its being in a reading frame other than the coding frame or to its being an intron sequence, a 3' or 5' untranslated sequence, or a regulatory sequence such as a promoter. A polypeptide also can be chemically synthesized and can be modified by chemical or enzymatic methods following translation or chemical synthesis. The terms "protein," "polypeptide" and "peptide" are used interchangeably herein when referring to a translated nucleic acid, for example, a gene product.

[0041] As used herein, the phrase "determining the identity of a target polypeptide" refers to determining at least one characteristic of the polypeptide, for example, the molecular mass or charge, or the identity of at least one amino acid, or identifying a particular pattern of peptide fragments of the target polypeptide. Determining the identity of a target polypeptide can be performed, for example, by using mass spectrometry to determine the amino acid sequence of at

least a portion of the polypeptide, or to determine the pattern of peptide fragments of the target polypeptide produced, for example, by treatment of the polypeptide with one or more endopeptidases.

[0042] In determining the identity of a target polypeptide, the number of nucleotide repeats encoding the target polypeptide can be quantified. As used herein, the term "quantify," when used in reference to nucleotide repeats encoding a target polypeptide, means a determination of the exact number of nucleotide repeats present in the nucleotide sequence encoding the target polypeptide. As disclosed herein, the number of nucleotide repeats, for example, trinucleotide repeats, can be quantified by using mass spectrometry to determine the number of amino acids, which are encoded by the repeat, that are present in the target polypeptide. It is recognized, however, that the number of nucleotide repeats encoding a target polypeptide need not be quantified to determine the identity of a target polypeptide, since a measure of the relative number of amino acids encoded by a region of nucleotide repeats also can be used to determine the identity of the target polypeptide by comparing the mass spectrum of the target polypeptide with that of a corresponding known polypeptide.

[0043] As used herein, the term "nucleotide repeats" refers to any nucleotide sequence containing tandemly repeated nucleotides. Such tandemly repeated nucleotides can be, for example, tandemly repeated dinucleotide, trinucleotide, tetranucleotide or pentanucleotide sequences, or any tandem array of repeated units.

[0044] As used herein, a reference polypeptide is a polypeptide to which the target polypeptide is compared in order to identify the polypeptide in methods that do not involve sequencing the polypeptide. Reference polypeptides typically are known polypeptides

[0045] As used herein, the term "conditioned" or "conditioning," when used in reference to a polypeptide, particularly a target polypeptide, means that the polypeptide is modified so as to decrease the laser energy required to volatilize the polypeptide, to minimize the likelihood of fragmentation of the polypeptide, or to increase the resolution of a mass spectrum of the polypeptide or of the component amino acids. Resolution of a mass spectrum of a target polypeptide can be increased by conditioning the polypeptide prior to performing mass spectrometry. Conditioning can be performed at any stage prior to mass spectrometry and, in particular, can be performed while the polypeptide is immobilized. A polypeptide can be conditioned, for example, by treating the polypeptide with a cation exchange material or an anion exchange material, which can reduce the charge heterogeneity of the polypeptide, thereby for eliminating peak broadening due to heterogeneity in the number of cations (or anions) bound to the various polypeptides in a population. Contacting a polypeptide with an alkylating agent such as alkyl iodide, iodoacetamide, iodoethanol, or 2,3-epoxy-1-propanol, the formation of disulfide bonds, for example, in a polypeptide can be prevented. Likewise, charged amino acid side chains can be converted to uncharged derivatives employing trialkylsilyl chlorides.

[0046] Conditioning of proteins is generally unnecessary because proteins are relatively stable under acidic, high energy conditions so that proteins do not require conditioning for mass spectrometric analyses. There are means of improving resolution, however, particularly for shorter peptides, such as by incorporating modified amino acids that are more basic than the corresponding unmodified residues. Such modification in general increases the stability of the polypeptide during mass spectrometric analysis. Also, cation exchange chromatography, as well as general washing and purification procedures which remove proteins and other reaction mixture components away from the target polypeptide, can be used to clean up the peptide after *in vitro* translation and thereby increase the resolution of the spectrum resulting from mass spectrometric analysis of the target polypeptide

[0047] As used herein, delayed extraction, refers to methods in which conditions are selected to permit a longer optimum extraction delay and hence a longer residence time, which results in increased resolution (see, *e.g.*, Juhasz et al. (1996) *Analysis, Anal. Chem.* 68:941-946; and Vestal et al. (1995) *Rapid Communications in Mass Spectrometry* 9:1044-1050; see also, *e.g.*, U.S. Patent No. 5,777,325, U.S. Patent No. 5,742,049, U.S. Patent No. 5,654,545, U.S. Patent No. 5,641,959, U.S. Patent No. 5,654,545 and U.S. Patent No. 5,760,393 for descriptions of MALDI and delayed extraction protocols). In particular, delayed ion extraction is a technique whereby a time delay is introduced between the formation of the ions and the application of the accelerating field. During the time lag, the ions move to new positions according to their initial velocities. By properly choosing the delay time and the electric fields in the acceleration region, the time of flight of the ions can be adjusted so as to render the flight time independent of the initial velocity to the first order. For example, a particular method involves exposure of the target polypeptide sample to an electric field before and during the ionization process, which results in a reduction of background signal due to the matrix, induces fast fragmentation and controls the transfer of energy prior to ion extraction.

[0048] As used herein, the term "multiplexing" refers to simultaneously determining the identity of at least two target polypeptides by mass spectrometry. For example, where a population of different target polypeptides are present in an array on a microchip or are present on another type of solid support, multiplexing can be used to determine the identity of a plurality of target polypeptides. Multiplexing can be performed, for example, by differentially mass modifying each different polypeptide of interest, then using mass spectrometry to determine the identity of each different polypeptide. Multiplexing provides the advantage that a plurality of target polypeptides can be identified in as few as a single mass spectrum, as compared to having to perform a separate mass spectrometry analysis for each individual target polypeptide

[0049] As used herein, the term "plurality," when used in reference to a polynucleotide or to a polypeptide, means two or more polynucleotides or polypeptides, each of which has a different nucleotide or amino acid sequence, respectively.

Such a difference can be due to a naturally occurring variation among the sequences, for example, to an allelic variation in a nucleotide or an encoded amino acid, or can be due to the introduction of particular modifications into various sequences, for example, the differential incorporation of mass modified amino acids into each polypeptide in a plurality.

[0050] As used herein, "*in vitro* transcription system" refers to a cell-free system containing an RNA polymerase and other factors and reagents necessary for transcription of a DNA molecule operably linked to a promoter that specifically binds an RNA polymerase. An *in vitro* transcription system can be a cell extract, for example, a eukaryotic cell extract. The term "transcription," as used herein, generally means the process by which the production of RNA molecules is initiated, elongated and terminated based on a DNA template. In addition, the process of "reverse transcription," which is well known in the art, is considered as encompassed within the meaning of the term "transcription" as used herein. Transcription is a polymerization reaction that is catalyzed by DNA-dependent or RNA-dependent RNA polymerases. Examples of RNA polymerases include the bacterial RNA polymerases, SP6 RNA polymerase, T3 RNA polymerase, T3 RNA polymerase, and T7 RNA polymerase

[0051] As used herein, the term "translation" describes the process by which the production of a polypeptide is initiated, elongated and terminated based on an RNA template. For a polypeptide to be produced from DNA, the DNA must be transcribed into RNA, then the RNA is translated due to the interaction of various cellular components into the polypeptide. In prokaryotic cells, transcription and translation are "coupled", meaning that RNA is translated into a polypeptide during the time that it is being transcribed from the DNA. In eukaryotic cells, including plant and animal cells, DNA is transcribed into RNA in the cell nucleus, then the RNA is processed into mRNA, which is transported to the cytoplasm, where it is translated into a polypeptide.

[0052] The term "translation system" refers to a cellular or cell-free system for performing a translation reaction. The term "cellular translation system" refers to a translation system based on a permeabilized cell; the term "cell-free translation system" or "*in vitro* translation system" refers to a cell extract or a reconstituted translation system. The term "reconstituted translation system" refers to a system containing purified or partially purified translation factors such as elongation factors. An *in vitro* translation system contains at least the minimum elements necessary for translation of an RNA molecule into a polypeptide. An *in vitro* translation system, which can be a eukaryotic or prokaryotic system, typically contains ribosomes, tRNA molecules, rRNA, an initiator methionyl-tRNA^{Met}, proteins or complexes involved in translation, for example, eukaryotic initiation factor 2 (eIF₂), eIF₃ and eIF_{4F}, and the cap-binding complex, including the cap-binding protein.

[0053] The term "isolated" as used herein with respect to a nucleic acid, including DNA and RNA, refers to nucleic acid molecules that are substantially separated from other macromolecules normally associated with the nucleic acid in its natural state. An isolated nucleic acid molecule is substantially separated from the cellular material normally associated with it in a cell or, as relevant, can be substantially separated from bacterial or viral material; or from culture medium when produced by recombinant DNA techniques; or from chemical precursors or other chemicals when the nucleic acid is chemically synthesized. In general, an isolated nucleic acid molecule is at least about 50% enriched with respect to its natural state, and generally is about 70% to about 80% enriched, particularly about 90% or 95% or more. Preferably, an isolated nucleic acid constitutes at least about 50% of a sample containing the nucleic acid, and can be at least about 70% or 80% of the material in a sample, particularly at least about 90% to 95% or greater of the sample. An isolated nucleic acid can be a nucleic acid fragment that does not occur in nature and, therefore, is not found in a natural state.

[0054] The term "isolated" also is used herein to refer to polypeptides that are substantially separated from other macromolecules normally associated with the polypeptide in its natural state. An isolated polypeptide can be identified based on its being enriched with respect to materials it naturally is associated with or its constituting a fraction of a sample containing the polypeptide to the same degree as defined above for an "isolated" nucleic acid, i.e., enriched at least about 50% with respect to its natural state or constituting at least about 50% of a sample containing the polypeptide. An isolated polypeptide, for example, can be purified from a cell that normally expresses the polypeptide or can be produced using recombinant DNA methodology.

[0055] As used herein, the term "nucleic acid" refers to a polynucleotide, including a deoxyribonucleic acid (DNA), a ribonucleic acid (RNA), and an analog of DNA or RNA containing, for example, a nucleotide analog or a "backbone" bond other than a phosphodiester bond; for example, a phosphotriester bond, a thioester bond, or a peptide bond (peptide nucleic acid). A nucleic acid can be single stranded or double stranded and can be, for example, a DNA-RNA hybrid. A nucleic acid also can be a portion of a longer nucleic acid molecule, for example, a portion of a gene containing a polymorphic region. The molecular structure of a nucleotide sequence, for example, a gene or a portion thereof, is defined by its nucleotide content, including deletions, substitutions or additions of one or more nucleotides; the nucleotide sequence; the state of methylation; or any other modification of the nucleotide sequence.

[0056] Reference to a nucleic acid as a "polynucleotide" is used in its broadest sense to mean two or more nucleotides or nucleotide analogs linked by a covalent bond, including single stranded or double stranded molecules. The term "oligonucleotide" also is used herein to mean two or more nucleotides or nucleotide analogs linked by a covalent bond, although those in the art will recognize that oligonucleotides such as PCR primers generally are less than about fifty to

one hundred nucleotides in length. The term "amplifying," when used in reference to a nucleic acid, means the repeated copying of a DNA sequence or an RNA sequence, through the use of specific or non-specific means, resulting in an increase in the amount of the specific DNA or RNA sequences intended to be copied.

[0057] A process as disclosed herein can be used to determine a nucleotide sequence of an unknown polynucleotide by comparing the amino acid sequence of a polypeptide encoded by the unknown polynucleotide with the amino acid sequence of a polypeptide encoded by a corresponding known polynucleotide. The determined nucleotide sequence of the unknown polynucleotide can be the same as a naturally occurring nucleotide sequence encoding the polypeptide, or can be different from the naturally occurring sequence due to degeneracy of the genetic code.

[0058] As used herein, the term "unknown polynucleotide" refers to a polynucleotide, the encoded polypeptide of which is being examined by mass spectrometry. Generally, an unknown polynucleotide is obtained from a biological sample. The term "corresponding known polynucleotide" means a defined counterpart of the unknown polynucleotide. A corresponding known polynucleotide generally is used as a control for comparison to the unknown polynucleotide and can be, for example, the nucleotide sequence of an allele of the unknown polynucleotide that is present in the majority of subjects in a population. For example, an "unknown polynucleotide" can be a DNA sequence that is obtained from a prostate cancer patient and includes the polymorphic region that demonstrates amplification of a trinucleotide sequence associated with prostate cancer, and the "corresponding known polynucleotide" can be the same polymorphic region from a subject that does not have prostate cancer, for example, from a female subject. An unknown polynucleotide also can be a mutated gene, which can alter the phenotype of a subject as compared to a subject not having the mutated gene. A mutated gene can be recessive, dominant or codominant, as is well known in the art.

[0059] The term "plasmid" refers generally to a circular DNA sequence which, in its vector form, is not bound to a chromosome. The terms "plasmid" and "vector" are used interchangeably herein, since the plasmid is the most commonly used form of a vector. Vectors such as a lambda vector can be linear but, nevertheless, are included within the meaning of the term "plasmid" or "vector" as used herein. Expression vectors and other vectors serving equivalent functions, and that become known in the art subsequently hereto, are included within the meaning of the term "plasmid" or "vector" as used herein.

[0060] In general, a nucleic acid encoding a polypeptide of interest, for example, a target polypeptide, is cloned into a plasmid and is operably linked to regulatory elements necessary for transcription or translation of the cloned nucleic acid. As used herein, the term "operably linked" means that a nucleic acid encoding a polypeptide is associated with a regulatory element, particularly a promoter, such that the regulatory element performs its function with respect to the nucleic acid molecule to which it is linked. For example, a promoter element that is operably linked to a nucleic acid allows for transcription of the nucleic acid when the construct is placed in conditions suitable for transcription to occur. It should be recognized that the term "regulatory element" is used broadly herein to include a nucleotide sequence, either DNA or RNA, that is required for transcription or translation, for example, a nucleotide sequence encoding a STOP codon or a ribosome binding site.

[0061] The term "target nucleic acid" refers to any nucleic acid of interest, including a portion of a larger nucleic acid such as a gene or an mRNA. A target nucleic acid can be a polymorphic region of a chromosomal nucleic acid, for example, a gene, or a region of a gene potentially having a mutation. Target nucleic acids include, but are not limited to, nucleotide sequence motifs or patterns specific to a particular disease and causative thereof, and to nucleotide sequences specific as a marker of a disease but not necessarily causative of the disease or condition. A target nucleic acid also can be a nucleotide sequence that is of interest for research purposes, but that may not have a direct connection to a disease or that may be associated with a disease or condition, although not yet proven so. A target nucleic acid can be any region of contiguous nucleotides that encodes a polypeptide of at least 2 amino acids, generally at least 3 or 4 amino acids, particularly at least 5 amino acids. A target nucleic acid encodes a target polypeptide.

[0062] The term "target polypeptide" refers to any polypeptide of interest that is subjected to mass spectrometry for the purposes disclosed herein, for example, for identifying the presence of a polymorphism or a mutation. A target polypeptide contains at least 2 amino acids, generally at least 3 or 4 amino acids, and particularly at least 5 amino acids. A target polypeptide can be encoded by a nucleotide sequence encoding a protein, which can be associated with a specific disease or condition, or a portion of a protein. A target polypeptide also can be encoded by a nucleotide sequence that normally does not encode a translated polypeptide. A target polypeptide can be encoded, for example, from a sequence of dinucleotide repeats or trinucleotide repeats or the like, which can be present in chromosomal nucleic acid, for example, a coding or a non-coding region of a gene, for example, in the telomeric region of a chromosome.

[0063] A process as disclosed herein also provides a means to identify a target polypeptide by mass spectrometric analysis of peptide fragments of the target polypeptide. As used herein, the term "peptide fragments of a target polypeptide" refers to cleavage fragments produced by specific chemical or enzymatic degradation of the polypeptide. The production of such peptide fragments of a target polypeptide is defined by the primary amino acid sequence of the polypeptide, since chemical and enzymatic cleavage occurs in a sequence specific manner. Peptide fragments of a target polypeptide can be produced, for example, by contacting the polypeptide, which can be immobilized to a solid support, with a chemical agent such as cyanogen bromide, which cleaves a polypeptide at methionine residues, or

hydroxylamine at high pH, which can cleave an Asp-Gly peptide bond; or with an endopeptidase such as trypsin, which cleaves a polypeptide at Lys or Arg residues.

[0064] The identity of a target polypeptide can be determined by comparison of the molecular mass or sequence with that of a reference or known polypeptide. For example, the mass spectra of the target and known polypeptides can be compared.

[0065] As used herein, the term "corresponding or known polypeptide" is a known polypeptide generally used as a control to determine, for example, whether a target polypeptide is an allelic variant of the corresponding known polypeptide. It should be recognized that a corresponding known protein can have substantially the same amino acid sequence as the target polypeptide, or can be substantially different. For example, where a target polypeptide is an allelic variant that differs from a corresponding known protein by a single amino acid difference, the amino acid sequences of the polypeptides will be the same except for the single difference. Where a mutation in a nucleic acid encoding the target polypeptide changes, for example, the reading frame of the encoding nucleic acid or introduces or deletes a STOP codon, the sequence of the target polypeptide can be substantially different from that of the corresponding known polypeptide.

[0066] As disclosed herein, a target polypeptide can be isolated using a reagent that interacts specifically with the target polypeptide, with a tag peptide fused to the target polypeptide, or with a tag conjugated to the target polypeptide. As used herein, the term "reagent" means a ligand or a ligand binding molecule that interacts specifically with a particular ligand binding molecule or ligand, respectively. The term "tag peptide" is used herein to mean a peptide that is specifically bound by a reagent. The term "tag" refers more generally to any molecule that is specifically bound by a reagent and, therefore, includes a tag peptide. A reagent can be, for example, an antibody that interacts specifically with an epitope of a target polypeptide or an epitope of a tag peptide. For example, a reagent can be an anti-myc epitope antibody, which can interact specifically with a myc epitope fused to a target polypeptide. A reagent also can be, for example, a metal ion such as nickel ion or cobalt ion, which interacts specifically with a polyhistidine tag peptide; or zinc, copper or, for example, a zinc finger domain, which interacts specifically with an polyarginine or polylysine tag peptide; or a molecule such as avidin, streptavidin or a derivative thereof, which interacts specifically with a tag such as biotin or a derivative thereof (see, e.g., U.S. application Serial No. 08/649,876, and also the corresponding published International PCT application No. WO 97/43617, which describe methods for dissociating biotin compounds, including biotin and biotin analogs conjugated (biotinylated) to the polypeptide, from biotin binding compounds, including avidin and streptavidin, using amines, particularly ammonia).

[0067] The term "interacts specifically," when used in reference to a reagent and the epitope, tag peptide or tag to which the reagent binds, indicates that binding occurs with relatively high affinity. As such, a reagent has an affinity of at least about $1 \times 10^6 \text{ M}^{-1}$, generally, at least about $1 \times 10^7 \text{ M}^{-1}$, and, in particular, at least about $1 \times 10^8 \text{ M}^{-1}$, for the particular epitope, tag peptide or tag. A reagent that interacts specifically, for example, with a particular tag peptide primarily binds the tag peptide, regardless of whether other unrelated molecules are present and, therefore, is useful for isolating the tag peptide, particularly a target polypeptide fused to the tag peptide, from a sample containing the target polypeptide, for example, from an *in vitro* translation reaction.

[0068] It can be advantageous in performing a disclosed process to conjugate a nucleic acid, for example, a target nucleic acid, or a polypeptide, for example, a target polypeptide, to a solid support such as a bead, microchip, glass or plastic capillary, or any surface, particularly a flat surface, which can contain a structure such as wells, pins or the like. A nucleic acid or a polypeptide can be conjugated to a solid support by various means, including, for example, by a streptavidin or avidin to biotin interaction; a hydrophobic interaction; by a magnetic interaction using, for example, functionalized magnetic beads such as DYNABEADS, which are streptavidin coated magnetic beads (Dynal Inc.; Great Neck NY; Oslo Norway); by a polar interaction such as a "wetting" association between two polar surfaces or between oligo/polyethylene glycol; by the formation of a covalent bond such as an amide bond, a disulfide bond, a thioether bond; through a crosslinking agent; and through an acid-labile or photocleavable linker (see, for example, Hermanson, "Bioconjugate Techniques" (Academic Press 1996)). In addition, a tag or a peptide such as a tag peptide can be conjugated to polypeptide of interest, particularly to a target polypeptide.

[0069] A process as disclosed herein is useful for determining the amino acid sequence of a polypeptide of interest, for example, by using an agent that cleaves amino acids from a terminus of the polypeptide to produce a nested set of deletion fragments of the polypeptide and cleaved amino acids, and using mass spectrometry to identify either the cleaved amino acids or the deletion fragments. As used herein, the phrase "agent that cleaves amino acids from a terminus of a polypeptide" refers to a means, which can be physical, chemical or biological, for removing a carboxyl terminal or an amino terminal amino acid from a polypeptide. A physical agent is exemplified by a light source, for example, a laser, that can cleave a terminal amino acid, particularly where the amino acid is bound to the polypeptide through a photolabile bond. A chemical agent is exemplified by phenylisothiocyanate (Edman's reagent), which, in the presence of an acid, cleaves an amino terminal amino acid from a polypeptide. A biological agent that cleaves an amino acid from a terminus of a polypeptide is exemplified by enzymes such as aminopeptidases and carboxypeptidases, which are well known in the art (see, for example, U.S. Patent No. 5,792,664; International Publ. No. WO 96/36732).

[0070] As used herein, the term "deletion fragment" refers to that portion of a polypeptide that remains following cleavage of one or more amino acids. The phrase "nested set of deletion fragments," when used in reference to a polypeptide to be sequenced, means a population of deletion fragments that results from sequential terminal cleavage of the amino acids of the polypeptide and that contains at least one deletion fragment that terminates in each amino acid of the portion of the polypeptide to be sequenced

[0071] A process as disclosed herein can be used to identify a subject that has or is predisposed to a disease or condition. As used herein, the term "disease" has its commonly understood meaning of a pathologic state in a subject. For purposes of the present disclosure, a disease can be due, for example, to a genetic mutation, a chromosomal defect or an infectious organism. The term "condition," which is to be distinguished from conditioning of a polypeptide, is used herein to mean any state of a subject, including, for example, a pathologic state or a state that determines, in part, how the subject will respond to a stimulus. The condition of a subject is determined, in part, by the subject's genotype, which can provide an indication as to how the subject will respond, for example, to a graft or to treatment with a particular medicament. Accordingly, reference to a subject being predisposed to a condition can indicate, for example, that the subject has a genotype indicating that the subject will not respond favorably to a particular medicament.

[0072] Reference herein to an allele or an allelic variant being "associated" with a disease or condition means that the particular genotype is characteristic, at least in part, of the genotype exhibited by a population of subjects that have or are predisposed to the disease or condition. For example, an allelic variant such as a mutation in the BRCA1 gene is associated with breast cancer, and an allelic variant such as a higher than normal number of trinucleotide repeats in a particular gene is associated with prostate cancer. The skilled artisan will recognize that an association of an allelic variant with a disease or condition can be identified using well known statistical methods for sampling and analysis of a population.

[0073] As used herein, the term "conjugated" refers to a stable attachment, which can be a covalent attachment or a noncovalent attachment, provided the noncovalent attachment is stable under the condition to which the bond is to be exposed. In particular, a polypeptide can be conjugated to a solid support through a linker, which can provide a non-cleavable, cleavable or reversible attachment.

[0074] As used herein, the term "solid support" means a flat surface or a surface with structures, to which a functional group, including a polypeptide containing a reactive group, can be conjugated. The term "surface with structures" is used herein to mean a support that contains, for example, wells, pins or the like, to which a functional group, including a polypeptide containing a reactive group, can be attached. Numerous examples of solid supports are disclosed herein or otherwise known in the art.

[0075] As used herein, the term "starting nucleic acid" refers to at least one molecule of a target nucleic acid, which encodes a target polypeptide. The starting nucleic acid can be DNA or RNA, including mRNA, and can be single stranded or double stranded, including a DNA-RNA hybrid. A mixture of any of these nucleic acids also can be employed as a starting nucleic acid for performing a process as disclosed herein, as can the nucleic acids produced following an amplification reaction.

[0076] It should be understood that the term "primer," as used herein, can refer to more than one primer, particularly in the case where there is some ambiguity in the information regarding the terminal sequence of a nucleic acid to be amplified. For example, where a nucleic acid sequence is inferred from protein sequence information, a collection of primers containing sequences representing all possible codon variations based on degeneracy of the genetic code is used for each strand. One primer from this collection is expected to be identical with a region of the sequence to be amplified.

[0077] A process is provided for determining the identity of a target polypeptide by using mass spectroscopy to determine the molecular mass of the target polypeptide and comparing it to the molecular mass of a polypeptide of known identity, thereby determining the identity of the target polypeptide. The identity of a target polypeptide can be, for example, the mass or amino acid sequence of at least a portion of the target polypeptide or by comparing the mass to a known polypeptide, which is a wild-type or known mutain.

[0078] A target polypeptide can be obtained from a subject, particularly from a cell or tissue in the subject or from a biological fluid. A target polypeptide also can be obtained by *in vitro* translation of an RNA molecule encoding the target polypeptide; or by *in vitro* transcription of a nucleic acid encoding the target polypeptide, followed by translation, which can be performed *in vitro* or in a cell, where the nucleic acid to be transcribed is obtained from a subject.

[0079] A process as disclosed herein provides a fast and reliable means for indirectly obtaining nucleic acid sequence information. Since the mass of a polypeptide is only about 10% of the mass of the corresponding DNA, the translated polypeptide generally is far more amenable to mass spectrometric detection than the corresponding nucleic acid. In addition, mass spectrometric detection of polypeptides yields analytical signals of far higher sensitivity and resolution than signals routinely obtained with DNA, due to the inherent instability of DNA to volatilization and its affinity for nonvolatile cationic impurities.

[0080] These processes are particularly useful for a number of applications, such as identifying mutations and thereby screening for certain genetic disorders. A process as disclosed herein also provides an efficient means for determining

the presence of a single base in a polynucleotide, for example, a single base mutation that introduces a STOP codon into an open reading frame of a gene, since such a mutation results in premature protein truncation; or a single base difference that results in a change in the encoded amino acid in an allelic variant of a polymorphic gene, since different amino acids can be distinguished based on their masses. Mutation screening by direct mass analysis of a gene such as p53 or BRCA1 requires a system that permits detection of a single base mutation, which can be difficult when examining a DNA sequence directly. A single base mutation resulting, for example, in a premature STOP codon, can radically change the mass of the encoded protein by truncation and, therefore, is readily identifiable using a process as disclosed herein. A single base change need not result in a STOP codon in order to be detectable, since a single base change that results in an amino acid change, for example, alanine to glycine, also is detectable using a process as disclosed herein (see Examples).

[0081] A process as disclosed herein can be used for identifying the presence of nucleotide repeats, particularly an abnormal number of nucleotide repeats, by determining the identity of a target polypeptide encoded by such repeats. As disclosed herein, an abnormal number of nucleotide repeats can be identified by using mass spectrometry to compare the mass of a target polypeptide with that of a corresponding known polypeptide.

[0082] In a particular application, the disclosed processes, can be used, for example, in detecting an abnormal number of CAG repeats in the SCA-1 gene or in detecting the presence of a nucleotide substitution from a C to a G in one of the trinucleotide repeats in a subject with spino-cerebellar ataxia 1 (SCA-1). Mass spectrometry is used to determine the molecular mass of a target polypeptide encoded by a nucleic acid containing the trinucleotide repeats and comparing the molecular mass of the target polypeptide with the molecular mass of a polypeptide encoded by a nucleic acid having a known number of trinucleotide repeats and a known nucleotide sequence (see Example 1). The identification of the nucleotide sequence of the target nucleic acid by this method is made possible, in part, due to the increased mass accuracy obtained by using mass spectrometry to detect the translation product, rather than directly detecting the nucleic acid by mass spectrometry.

[0083] For illustrative purposes, the open reading frame of the gene containing the (CAG)_x repeat associated with SCA-1 is shown in Figure 1. The SCA-1 sequence contains, in addition to a nonvariable stretch of 12 CAG repeats, a variable stretch that is shown in Figure 1A as containing 10 CAG repeats. As shown in Figure 1A, the SCA-1 gene encodes a 7.5 kiloDalton (kDa) protein containing 10 consecutive glutamine (Q) residues (Figure 1B). Accurate direct mass analysis of the 60 kDa 200-mer shown in Figure A with currently available mass spectrometric instrumentation would be challenging. A recent study of the SCA-1 gene showed that 25 to 36 repeat units generally are present in unaffected subjects, while affected subjects have 43 to 81 repeat units. Assuming a worst case of 81 repeat units, 213 bases in addition to the 200-mer shown in Figure 1A would have to be detected with sufficient resolution. A nucleotide sequence of greater than about a 400-mer (> 120 kDa) has not been detected satisfactorily by mass spectrometry. In comparison, analysis of the translation product for the sequence having 81 repeats requires mass measurement of only about 137 amino acid residues (about 15 kDa). A typical 0.3% mass accuracy for low resolution instrumentation results in a maximum 13 Dalton error, which is far lower than the mass of a single amino acid residue. Accordingly, far better than single amino acid resolution can be obtained with a process for determining the identity of a target polypeptide as disclosed herein.

OBTAINING A TARGET POLYPEPTIDE

[0084] Any polypeptide for which identifying information is required is contemplated herein as a target polypeptide. The polypeptide may be obtained from any source. A target polypeptide, or a target nucleic acid encoding the polypeptide, is can be obtained from a subject, which is typically a mammal, particularly a human. Generally, the target polypeptide is isolated prior to mass spectrometry so as to permit the determination of the molecular mass of the polypeptide by mass spectrometric analysis. The degree of to which a polypeptide must be isolated for mass spectrometry is known in the art and varies depending on the type of mass spectrometric analysis performed.

[0085] A target polypeptide can be a portion of a protein, and can be obtained using methods known in the art. For example, a protein can be isolated from a biological sample using an antibody, then can be cleaved using a proteinase that cuts selectively at specific amino acid sequences, and the target polypeptide can be purified by a method such as chromatography or electrophoresis. Thus, a process as disclosed herein can be performed, for example, by subjecting a protein, which contains a target polypeptide, to limited proteolysis; isolating the target polypeptide; and examining it by mass spectrometric analysis, thereby providing a means for determining the identity of the target polypeptide.

[0086] An antibody, or antigen binding fragment of an antibody, that interacts specifically with an epitope present on a polypeptide of interest is characterized by having specific binding activity for the epitope of at least about $1 \times 10^6 \text{ M}^{-1}$, generally, at least about $1 \times 10^7 \text{ M}^{-1}$ or greater. Accordingly, Fab, F(ab')₂, Fd and Fv fragments of an antibody that retain specific binding activity for a particular epitope are included within the meaning of the term antibody.

[0087] An antibody useful for isolating a polypeptide of interest, particularly a target polypeptide, can be a naturally occurring antibody or a non-naturally occurring antibody, including, for example, a single chain antibody, a chimeric

antibody, a bifunctional antibody or a humanized antibody, as well as an antigen-binding fragment of such antibodies. Such non-naturally occurring antibodies can be constructed using solid phase peptide synthesis, can be produced recombinantly or can be obtained, for example, by screening combinatorial libraries containing of variable heavy chains and variable light chains (see Huse et al., Science 246:1275-1281 (1989)). These and other methods of making, for

5 example, chimeric, humanized, CDR-grafted, single chain, and bifunctional antibodies are well known to those skilled in the art (Winter and Harris, Immunol. Today 14:243-246 (1993); Ward et al., Nature 341:544-546 (1989); Hilyard et al., Protein Engineering: A practical approach (IRL Press 1992); Borrabeck, Antibody Engineering, 2d ed. (Oxford University Press 1995); Harlow and Lane, "Antibodies: A laboratory manual" (Cold Spring Harbor Laboratory Press 1988)).

10 [0088] An antibody useful for isolating a target polypeptide can be obtained from a commercial source, or can be raised using a protein containing the target polypeptide, or a peptide portion thereof, as an immunogen, or using an epitope that is fused to the polypeptide, for example, a myc epitope. Such an immunogen can be prepared from natural sources or produced recombinantly, or can be synthesized using routine chemical methods. An otherwise non-immunogenic epitope can be made immunogenic by coupling the hapten to a carrier molecule such bovine serum albumin (BSA) or keyhole limpet hemocyanin (KLH), or by expressing the epitope as a fusion protein. Various other carrier

15 molecules and methods for coupling a hapten to a carrier molecule are well known in the art (see, for example, Harlow and Lane, "Antibodies: A laboratory manual" (Cold Spring Harbor Laboratory Press 1988)).

[0089] An antibody that interacts specifically with a polypeptide of interest, particularly a target polypeptide or peptide portion thereof, is useful, for example, for determining whether the target polypeptide is present in a biological sample. The identification of the presence or level of the target polypeptide can be made using well known immunoassay and

20 immunohisto-chemical methods (Harlow and Lane, "Antibodies: A laboratory manual" (Cold Spring Harbor Laboratory Press 1988)). In particular, an antibody that interacts specifically with a tag peptide fused to a target polypeptide can be used to isolate the target polypeptide from a sample, which can be, for example, a biological sample or an *in vitro* translation reaction.

[0090] Methods for raising polyclonal antibodies, for example, in a rabbit, goat, mouse or other mammal, are well known in the art (Harlow and Lane, "Antibodies: A laboratory manual" (Cold Spring Harbor Laboratory Press 1988)). In addition, monoclonal antibodies can be obtained using methods that are well known and routine in the art (Harlow and Lane, "Antibodies: A laboratory manual" (Cold Spring Harbor Laboratory Press 1988)). Essentially, spleen cells from a mouse immunized with a polypeptide of interest, or a peptide portion thereof, can be fused to an appropriate myeloma cell line such as SP/02 myeloma cells to produce hybridoma cells. Cloned hybridoma cell lines can be screened using

30 the immunizing polypeptide to identify clones that secrete appropriately specific antibodies. Hybridomas expressing antibodies having a desirable specificity and affinity can be isolated and utilized as a continuous source of the antibodies. Similarly, a recombinant phage that expresses, for example, a single chain antibody of interest also provides a monoclonal antibody.

[0091] Isolation and identification of a target polypeptide can be facilitated by linking a tag to the polypeptide, for example, by fusing the polypeptide to a tag peptide. Such a fusion polypeptide can be obtained, for example, by *in vitro* transcription and translation of a nucleotide sequence encoding the target polypeptide linked in frame to a nucleotide sequence encoding the tag peptide, then isolating the fusion polypeptide from the translation reaction using a reagent that interacts specifically with the tag peptide. The tag peptide can be, for example, a myc epitope or a peptide portion of the *Haemophilus influenza* hemagglutinin protein, against which specific antibodies can be prepared and also are

40 commercially available. A tag peptide also can be a polyhistidine sequence, for example, a hexahistidine sequence (His-6), which interacts specifically with metal ions such as zinc, nickel, or cobalt ions, or a polylysine or polyarginine sequence, comprising at least about four lysine or four arginine residues, respectively, which interact specifically with zinc, copper or, for example a zinc finger protein.

[0092] A tag can be also can be added to the polypeptide either by chemical modification of the polypeptide during or following its synthesis. For example, a target polypeptide containing a tag can be obtained by isolation from an *in vitro* translation reaction of a target nucleic acid molecule, where the translation reaction is performed in the presence of a modified amino acid and, if appropriate, a mis-aminoacylated tRNA carrying the modified amino acid. The modification of the amino acid is selected so that it contains a tag that allows the isolation of a polypeptide containing the modified amino acid. For example, a lysine residue can be replaced with a biotinylated lysine analog (or other lysine analog containing a tag) in the translation reaction, resulting in a translated polypeptide that contains biotinylated lysine residues. Such a tagged polypeptide can be isolated by affinity chromatography on a bed of immobilized avidin or streptavidin, for example. Other modified amino acids are disclosed in the U.S. Patent No. 5,643,722.

50 [0093] A target polypeptide can be isolated by affinity purification using, for example, an antibody, avidin or other specific reagent linked to a solid support. In such a method, the translation reaction is poured over the support, which can be present, for example, in a column, and the polypeptide is bound due to its specifically interacting with the reagent. For example, a target polypeptide fused to a polyhistidine tag peptide can be isolated on a column or bed of chelated nickel ions, whereas a target polypeptide fused to a polylysine or polyarginine tag can be isolated on a column or bed of chelated zinc or copper ions. Beds or columns having such divalent metal ions chelated thereto can be obtained from

a commercial source or prepared using methods known in the art. The polypeptide then can be eluted from the column in an isolated form and subjected to mass spectrometry.

ISOLATION OF A NUCLEIC ACID ENCODING A TARGET POLYPEPTIDE

[0094] In other embodiments, the polypeptide may be prepared from nucleic acid that encodes it. Thus, the target polypeptide can be isolated from a cell or tissue of the subject; or can be synthesized *in vitro* from an RNA molecule, for example, by *in vitro* translation, or from a DNA molecule by *in vitro* transcription and translation; or can be synthesized in a eukaryotic or prokaryotic host cell that is transformed with a target nucleic acid, which encodes the target polypeptide.

[0095] In preferred embodiments herein, a target polypeptide is isolated from a cell, a tissue or an *in vitro* translation system, for example, a reticulocyte lysate system. *In vitro* translation or *in vitro* transcription followed by translation are among the preferred methods of preparation of the polypeptides. The polypeptides can be purified after translation using any method known to those of skill in the art for purification. For example, the polypeptide can be isolated using a reagent that interacts specifically with the target polypeptide or with a protein containing the target polypeptide. Such a reagent can be an antibody that interacts specifically with an epitope of the target polypeptide, for example, an antibody to an epitope encoded by a trinucleotide repeat sequence. If the target polypeptide contains an amino acid that can be any of several amino acids, for example, where the target polypeptide is from a mutated protein, the antibody preferably interacts with an epitope that does not include an epitope containing the mutated amino acid(s). Antibodies that interact specifically with a protein containing a target polypeptide, or with the target polypeptide, can be prepared using methods well known in the art (Harlow and Lane, "Antibodies: A laboratory manual" (Cold Spring Harbor Laboratory Press 1988)).

[0096] A target polypeptide can be obtained from an RNA molecule, for example, by *in vitro* translation of the RNA molecule. The target polypeptide also can be obtained from a DNA molecule, where *in vitro* transcription of at least a portion of the DNA molecule is performed prior to translation. In particular, at least a portion of the DNA molecule containing the nucleotide sequence encoding the target polypeptide can be amplified, for example, by PCR prior to performing *in vitro* transcription or translation. Accordingly, a process for determining the identity of a target polypeptide, as disclosed herein, can include a step of isolating a target nucleic acid molecule, which can be DNA or RNA and from which the target polypeptide is obtained.

[0097] A nucleic acid sample, in an isolated or unisolated form, can be utilized as a starting nucleic acid in a method as disclosed herein, provided the sample is suspected of containing the target nucleic acid. The target nucleic acid can be a portion of a larger molecule or can be present initially as a discrete molecule such that the specific sequence constitutes the entire nucleic acid.

[0098] It is not necessary that a starting nucleic acid contain only the target nucleic acid in an isolated form. Provided that the starting nucleic acid is in an isolated form, the target nucleic acid can be a minor fraction of a complex mixture, for example, a portion of the β -globin gene contained in whole human DNA, or a portion of nucleic acid sequence of a particular microorganism that constitutes only a minor fraction of a particular biological sample. A starting nucleic acid also can contain more than one population of target nucleic acids.

[0099] The starting nucleic acid can be obtained from any source, including a natural source such as bacteria, yeast, viruses, protists, and higher organisms, including plants or animals, particularly from tissues, cells or organelles of such sources, or can be obtained from a plasmid such as pBR322, in which the nucleic acid previously was cloned. The starting nucleic acid can represent a sample of DNA, for example, isolated from an animal, particularly a mammal such as a human subject, and can be obtained from any cell source or body fluid. Examples of cell sources available in clinical practice include, but are not limited to, blood cells, buccal cells, cervico-vaginal cells, epithelial cells from urine, or cells present in a tissue obtained, for example, by biopsy. Body fluids include blood, urine and cerebrospinal fluid, as well as tissue exudates from a site of infection or inflammation.

[0100] A nucleic acid molecule can be extracted from a cell source or body fluid using any of numerous methods well known and routine in the art, and the particular method used to extract the nucleic acid will be selected as appropriate for the particular biological sample, including whether the nucleic acid to be isolated is DNA or RNA (see, for example, Sambrook et al., Molecular Cloning: A laboratory manual (Cold Spring Harbor Laboratory Press 1989)). For example, freeze-thaw and alkaline lysis procedures can be useful for obtaining nucleic acid molecules from solid materials such as cell or tissue samples; heat and alkaline lysis procedures can be useful for obtaining nucleic acid molecules from urine; and proteinase K extraction or phenol extraction can be useful to obtain nucleic acid from cells or tissues such as a blood sample (Rolff et al., "PCR: Clinical diagnostics and research" (Springer Verlag Publ. 1994)).

[0101] For utilization of a target nucleic acid from cells, the cells can be suspended in a hypotonic buffer and heated to about 90°C to 100°C for about 1 to 15 minutes, until cell lysis and dispersion of intracellular components occur. After the heating step, amplification reagents, if desired, can be added directly to the lysate. Such a direct amplification method can be used, for example, on peripheral blood lymphocytes or amniocytes. The amount of DNA extracted for analysis of human genomic DNA generally is at least about 5 pg, which corresponds to about 1 cell equivalent of a genome size of 4×10^9 base pairs. In some applications, for example, detection of sequence alterations in the genome of a microor-

ganism, variable amounts of DNA can be extracted.

[0102] In general, the nucleotides forming a polynucleotide are naturally occurring deoxyribonucleotides, such as adenine, cytosine, guanine or thymine linked to 2'-deoxyribose, or ribonucleotides such as adenine, cytosine, guanine or uracil linked to ribose. A polynucleotide also includes nucleotide analogs, including non-naturally occurring synthetic nucleotides or modified naturally occurring nucleotides. Such nucleotide analogs are well known in the art and are commercially available, as are polynucleotides containing such nucleotide analogs (Lin et al., Nucl. Acids Res. 22: 5220-5234 (1994); Jellinek et al., Biochemistry 34:11363-11372 (1995); Pagratis et al., Nature Biotechnol. 15:68-73 (1997)). The covalent bond linking the nucleotides of a polynucleotide generally is a phosphodiester bond. The covalent bond also can be any of numerous other bonds, including a thiodiester bond, a phosphorothioate bond, a peptide-like bond or any other bond known to those in the art as useful for linking nucleotides to produce synthetic polynucleotides (see, for example, Tam et al., Nucl. Acids Res. 22:977-986 (1994); Ecker and Crooke, BioTechnology 13:351360 (1995)).

[0103] Where it is desired to synthesize a polynucleotide for use in a process as disclosed herein the artisan will know that the selection of particular nucleotides or nucleotide analogs and the covalent bond used to link the nucleotides will depend, in part, on the purpose for which the polynucleotide is prepared. For example, where a polynucleotide will be exposed to an environment containing substantial nuclease activity, the artisan will select nucleotide analogs or covalent bonds that are relatively resistant to the nucleases. A polynucleotide containing naturally occurring nucleotides and phosphodiester bonds can be chemically synthesized or can be produced using recombinant DNA methods, using an appropriate polynucleotide as a template. In comparison, a polynucleotide containing nucleotide analogs or covalent bonds other than phosphodiester bonds generally will be chemically synthesized, although an enzyme such as T7 polymerase can incorporate certain types of nucleotide analogs and, therefore, can be used to produce such a polynucleotide recombinantly from an appropriate template (Jellinek et al., Biochemistry 34:11363-11372 (1995)).

[0104] A polynucleotide, for example, an oligonucleotide, that specifically hybridizes to a nucleic acid, particularly to a target nucleic acid or to sequences flanking a target nucleic acid is particularly useful. Such a hybridizing polynucleotide is characterized, in part, in that it is at least nine nucleotides in length, such sequences being particularly useful as primers for the polymerase chain reaction (PCR), and can be at least fourteen nucleotides in length or, if desired, at least seventeen nucleotides in length, such nucleotide sequences being particularly useful as hybridization probes, as well as for PCR. It should be recognized that the conditions required for specific hybridization of a first polynucleotide, for example, a PCR primer, with a second polynucleotide, for example, a target nucleic acid, depends, in part, on the degree of complementarity shared between the sequences, the GC content of the hybridizing molecules, and the length of the antisense nucleic acid sequence, and that conditions suitable for obtaining specific hybridization can be calculated based on readily available formulas or can be determined empirically (Sambrook et al., Molecular Cloning: A laboratory manual (Cold Spring Harbor Laboratory Press 1989; Ausubel et al., Current Protocols in Molecular Biology (Green Publ., NY 1989)).

TRANSCRIPTION AND TRANSLATION OF A TARGET NUCLEIC ACID

[0105] A target polypeptide can be obtained by translating an RNA molecule encoding the target polypeptide *in vitro*. If desired, the RNA molecule can be obtained by *in vitro* transcription of a nucleic acid, generally DNA, encoding the target polypeptide. Translation of a target polypeptide can be effected by directly introducing an RNA molecule encoding the polypeptide into an *in vitro* translation reaction or by introducing a DNA molecule encoding the polypeptide into an *in vitro* transcription/translation reaction or into an *in vitro* transcription reaction, then transferring the RNA to an *in vitro* translation reaction.

[0106] For *in vitro* transcription, the target DNA is operably linked to a promoter, from which transcription is initiated in the presence of an RNA polymerase capable of interacting with the promoter, ribonucleotides, and other reagents necessary for *in vitro* transcription. *In vitro* transcription can be performed as a separate step from an *in vitro* translation reaction or can be carried out in a single reaction, using well known methods (see, for example, Sambrook et al., Molecular Cloning: A laboratory manual (Cold Spring Harbor Laboratory Press 1989; see, also, U.S. Patent No. 4,766,072, which describes vectors useful for *in vitro* transcription). *In vitro* transcription kits are well known and are commercially available (Promega Corp., Madison WI).

[0107] An *in vitro* transcription reaction is carried out by incubating a template DNA, which generally includes the target nucleic acid, for about 1 hour at 37°C or 40°C, depending on the polymerase, in the presence of ribonucleotides, a cap analog such as GpppG or a methylated derivative thereof, an RNAase inhibitor, an RNA polymerase that recognizes the promoter operably linked upstream of the DNA to be transcribed, and an appropriate buffer containing Tris-HCl, MgCl₂, spermidine and NaCl. Following the transcription reaction, RNAase-free DNase can be added to remove the DNA template and the RNA purified, for example, by phenol-chloroform extraction (see, Sambrook et al., Molecular Cloning: A laboratory manual (Cold Spring Harbor Laboratory Press 1989). Usually about 5 to 10 µg of RNA is obtained per microgram of template DNA.

[0108] Where RNA is produced in a prokaryotic *in vitro* transcription system, the RNA can be produced in an uncapped

form, such as by *in vitro* transcription in the absence of a cap analog, since translation of RNA in a prokaryotic system does not require the presence of a cap such as N₇-methyl-G covalently linked to the 5' end of the mRNA. Capped RNA is translated much more efficiently than uncapped RNA in eukaryotic systems and, therefore, it can be desirable to cap the RNA during transcription or during translation when using a eukaryotic translation system. The *in vitro* transcribed RNA can be isolated, for example, by ethanol precipitation, then used for *in vitro* translation.

[0109] Translation systems can be cellular or cell-free and can be prokaryotic or eukaryotic. Cellular translation systems generally utilize intact cells, for example, oocytes, or utilize permeabilized cells, whereas cell-free (*in vitro*) translation systems utilize cell or tissue lysates or extracts, purified or partially purified components, or combinations thereof.

[0110] *In vitro* translation systems are well known and are commercially available and many different types and systems are well known and routinely used. Examples of *in vitro* translation systems include eukaryotic cell lysates such as rabbit reticulocyte lysates, rabbit oocyte lysates, human cell lysates, insect cell lysates and wheat germ extracts. Such lysates and extracts are can be prepared or are commercially available (Promega Corp.; Stratagene, La Jolla CA; Amersham, Arlington Heights IL; and GIBCO/BRL, Grand Island NY). *In vitro* translation systems generally contain macromolecules such as enzymes; translation, initiation and elongation factors; chemical reagents; and ribosomes. Mixtures of purified translation factors, as well as combinations of lysates or lysates supplemented with purified translation factors such as initiation factor-1 (IF-1), IF-2, IF-3 (alpha or beta), elongation factor T (EF-Tu) or termination factors, also can be used for mRNA translation *in vitro*.

[0111] Incubation times for *in vitro* translation range from about 5 minutes to many hours, but generally are about thirty minutes to five hours, usually about one to three hours. Incubation can be performed in a continuous manner, whereby reagents are flowed into the system and nascent polypeptides removed or left to accumulate, using a continuous flow system as described by Spirin et al. (Science 242:1162-64 (1988)). Such a process can be desirable for large scale production of nascent polypeptides. Incubation times vary significantly with the volume of the translation mix and the temperature of the incubation. Incubation temperatures can be between about 4°C to 60°C, generally about 15°C to 50°C, and usually about 25°C to 45°C, particularly about 25°C or about 37°C.

[0112] Translation reactions generally contain a buffer such as Tris-HCl, HEPES, or other suitable buffering agent to maintain the solution at about pH 6 to pH 8, generally about pH 7. Other components of a translation system can include dithiothreitol (DTT) or 2-mercaptoethanol as reducing agents, RNasin to inhibit RNA breakdown, and nucleoside triphosphates or creatine phosphate and creatine kinase to provide chemical energy for the translation process.

[0113] An *in vitro* translation system can be a reticulocyte lysate, which is available commercially or can be prepared according to methods disclosed herein or otherwise known in the art. Commercially available reticulocyte lysates are available, for example, from New England Nuclear and Promega Corp. (Cat. #L4960, L4970, and L4980). An *in vitro* translation system also can be a wheat germ translation system, which is available commercially or can be prepared according to well known methods. Commercially available wheat germ extracts can be obtained, for example, from Promega Corp. (for example, Cat # L4370). An *in vitro* translation system also can be a mixture of a reticulocyte lysate and a wheat germ extract, as can be obtained commercially (for example, Promega Corp., catalog # L4340). Other useful *in vitro* translation systems include *E. coli* extracts, insect cell extracts and frog oocyte extracts.

[0114] A rabbit reticulocyte lysate can be prepared as follows. Rabbits are rendered anemic by inoculation with acetylphenylhydrazine. About 7 days later, the rabbits are bled and the blood is collected and mixed with an ice cold salt solution containing NaCl, magnesium acetate (MgAc), KCl, and heparin. The blood mixture is filtered through a cheesecloth, centrifuged, and the buffy coat of white cells is removed. The pellet, which contains erythrocytes and reticulocytes, is washed with the salt solution, then lysed by the addition of an equal volume of cold water. Endogenous RNA is degraded by treating the lysate with micrococcal nuclease and calcium ions, which are necessary for nuclease activity, and the reaction is stopped by the addition of EGTA, which chelates the calcium ions and inactivates the nuclease. Hemin (about 20 to 80 µM), which is a powerful suppressor of an inhibitor of the initiation factor eIF-2, also can be added to the lysate. Translation activity of the lysates can be optimized by the addition of an energy generating system, for example, phosphocreatine kinase and phosphocreatine. The lysates then can be aliquoted and stored at -70°C or in liquid nitrogen. Further details regarding such a protocol are known (see, e.g., Sambrook et al., Molecular Cloning: A laboratory manual (Cold Spring Harbor Laboratory Press 1989).

[0115] An *in vitro* translation reaction using a reticulocyte lysate can be carried out as follows. Ten µl of a reticulocyte lysate, which can be prepared as disclosed above or can be obtained commercially, is mixed with spermidine, creatine phosphate, amino acids, HEPES buffer (pH 7.4), KCl, MgAc and the RNA to be translated, and incubated for an appropriate time, generally about one hour at 30°C. The optimum amount of MgAc for obtaining efficient translation varies from one reticulocyte lysate preparation to another and can be determined using a standard preparation of RNA and a concentration of MgAc varying from 0 to 1 mM. The optimal concentration of KCl also can vary depending on the specific reaction. For example, 70 mM KCl generally is optimal for translation of capped RNA, whereas 40 mM generally is optimal for translation of uncapped RNA. Optionally, the translation process is monitored by a method such as mass spectrometric analysis. Monitoring also can be performed, for example, by adding one or more radioactive amino acids such as ³⁵S-methionine and measuring incorporation of the radiolabel into the translation products by precipitating the

proteins in the lysate such as with TCA and counting the amount of radioactivity present in the precipitate at various times during incubation. The translation products also can be analyzed by immunoprecipitation or by SDS-polyacrylamide gel electrophoresis (see, for example, Sambrook et al., Molecular Cloning: A laboratory manual (Cold Spring Harbor Laboratory Press 1989; Harlow and Lane, "Antibodies: A laboratory manual" (Cold Spring Harbor Laboratory Press 1988)).

[0116] A wheat germ extract can be prepared as described by Roberts and Paterson (Proc. Natl. Acad. Sci., USA 70: 2330-2334 (1973)) and can be modified as described by Anderson (Meth. Enzymol. 101:635 (1983)), if desired. The protocol also can be modified according to manufacturing protocol L418 (Promega Corp.). Generally, wheat germ extract is prepared by grinding wheat germ in an extraction buffer, followed by centrifugation to remove cell debris. The supernatant is separated by chromatography from endogenous amino acids and from plant pigments that are inhibitory to translation. The extract also is treated with micrococcal nuclease to destroy endogenous mRNA, thereby reducing background translation to a minimum. The wheat germ extract contains the cellular components necessary for protein synthesis, including tRNA, rRNA and initiation, elongation and termination factors. The extract can be optimized further by the adding an energy generating system such as phosphocreatine kinase and phosphocreatine; MgAc is added at a level recommended for the translation of most mRNA species, generally about 6.0 to 7.5 mM magnesium.

[0117] *In vitro* translation in wheat germ extracts can be performed as described, for example, Erickson and Blobel (Meth. Enzymol. 96:38 (1982)), and can be modified, for example, by adjusting the final ion concentrations to 2.6 mM magnesium and 140 mM potassium, and the pH to 7.5 (U.S. Patent No. 4,983,521). Reaction mixtures can be incubated at 24°C for 60 minutes. Translations in wheat germ extracts can also be performed as described in U.S. Patent No. 5,492,817

[0118] *In vitro* translation reactions can be optimized by the addition of ions or other reagents. For example, magnesium is important for optimal translation, as it enhances the stability of assembled ribosomes and functions in their binding together during translation. Magnesium also appears to facilitate polymerase binding. Potassium also is important for optimizing translation but, unlike magnesium, for coupled transcription and translation reactions, the potassium ion concentration need not be altered beyond standard translation preparation levels.

[0119] Potassium and magnesium are in the standard rabbit reticulocyte lysate and their levels are partially from the endogenous lysate level and partially from the additions made in the preparation of the lysate, as are done for translation lysates. Since the magnesium concentration should be adjusted within a rather narrow range for optimal translation, the lysate magnesium levels should be measured directly through the use of a magnesium assay, prior to the addition of extra magnesium, so that the amount of magnesium in a reaction can be standardized from one batch of lysate to the next. The Lancer "Magnesium Rapid Stat Diagnostic Kit" (Oxford Lab Ware Division, Sherwood Medical Co.; St. Louis MO) is a useful assay for accurately measuring the magnesium level in a biological fluid. Once the magnesium concentration for a given batch of lysate is determined, additional magnesium, for example, in the form of a concentrated magnesium salt solution, can be added in a known manner to bring the magnesium concentration of the lysate to within the optimal range or, in the case of a modified lysate preparation to be used as one-half of a reaction mixture, to within twice the optimal range. The final magnesium concentration of rabbit reticulocyte lysate is adjusted, for example, by adding a concentrated solution of MgCl₂ or MgAc to a concentration greater than 2.5 mM, but less than 3.5 mM, generally between 2.6 mM and 3.0 mM.

[0120] A common addition to an *in vitro* translation reaction is an amount of a polyamine sufficient to stimulate efficient chain elongation. Accordingly, spermidine can be added to a reticulocyte lysate translation reaction to a final concentration of about 0.2 mM. Spermidine also can be added to wheat germ extracts, generally at a concentration of about 0.9 mM. Since the presence of polyamines lowers the effective magnesium concentration in a reaction, the presence of spermidine in a translation reaction should be considered when determining the appropriate concentration of magnesium to use. DTT also is added to the translation mixture, generally at a final concentration of about 1.45 mM in reticulocyte lysates and about 5.1 mM in wheat germ extracts.

[0121] Translation systems can be supplemented with additional factors such as tRNA molecules, which are commercially available (Sigma Chemical, St. Louis MO; Promega Corp., Madison WI; Boehringer Mannheim Biochemicals, Indianapolis IN) or can be prepared from *E. coli*, yeast, calf liver or wheat germ using well known methods. Isolation and purification of tRNA molecules involve cell lysis and phenol extraction, followed by chromatography on DEAE-cellulose. Amino acid-specific tRNA, for example, tRNA ^{fMet}, can be isolated by expression from cloned genes and overexpressed in host cells and separated from total tRNA in high yield and purity using, for example, preparative polyacrylamide gel electrophoresis, followed by band excision and elution (Seong and RajBhandary, Proc. Natl. Acad. Sci., USA 84: 334-338, 1987)).

[0122] Translation efficiency can be improved by adding RNAase inhibitors such as RNASIN or heparin to the translation reaction. RNASIN can be obtained, for example, from Promega Corp. (Cat # N2514). About 40 units of RNASIN are added to a 50 µl reaction. Although the addition of an RNAase inhibitor to reticulocyte lysates is not crucial, only limited translation occurs if an RNAase inhibitor is not added to a wheat germ extract translation reaction.

[0123] The translation process, including the movement of the ribosomes on the RNA molecules, is inhibited at an

appropriate time by the addition of an inhibitor of translation, for example, cycloheximide at a final concentration of 1 $\mu\text{g/ml}$. Magnesium ion, for example, MgCl_2 , at a concentration of about 5 mM also can be added to maintain the mRNA-80S ribosome-nascent polypeptide complexes (polysomes).

[0124] For determining the optimal *in vitro* translation conditions, translation of mRNA in an *in vitro* system can be monitored, for example, by mass spectrometric analysis. Alternatively, a labeled amino acid such as ^{35}S -methionine can be included in the translation reaction together with an amino acid mixture lacking this specific amino acid (e.g., methionine). A labeled non-radioactive amino acid also can be incorporated into a nascent polypeptide. For example, the translation reaction can contain a mis-aminoacylated tRNA (U.S. Patent No. 5,643,722). For example, a non-radioactive marker can be mis-aminoacylated to a tRNA molecule and the tRNA amino acid complex is added to the translation system. The system is incubated to incorporate the non-radioactive marker into the nascent polypeptide and polypeptides containing the marker can be detected using a detection method appropriate for the marker. Mis-aminoacylation of a tRNA molecule also can be used to add a marker to the polypeptide in order to facilitate isolation of the polypeptide. Such markers include, for example, biotin, streptavidin and derivatives thereof (see U.S. Patent No. 5,643,722). The translation process can also be followed by mass spectrometric analysis, which does not require the use of radioactivity or other label.

[0125] *In vitro* transcription and translation reactions can be performed simultaneously using, for example, a commercially available system such as the Coupled Transcription/Translation System (Promega Corp, catalog # L4606, # 4610 or # 4950). Coupled transcription and translation systems using RNA polymerases and eukaryotic lysates are described in U.S. Patent No. 5,324,637. Coupled *in vitro* transcription and translation also can be carried out using a prokaryotic system such as a bacterial system, for example, *E. coli* S30 cell-free extracts (Zubay, Ann. Rev. Genet. 7:267 (1973)). Although such prokaryotic systems allow coupled *in vitro* transcription and translation, they also can be used for *in vitro* translation only. When using a prokaryotic translation system, the RNA should contain sequence elements necessary for translation of an RNA in a prokaryotic system. For example, the RNA should contain prokaryotic ribosome binding sites, which can be incorporated into a target nucleic acid sequence during amplification using a primer containing the prokaryotic ribosome binding sequence. The ribosome binding sequence is positioned downstream of a promoter for use in *in vitro* transcription.

[0126] Cellular translation systems can be prepared as follows. Cells are permeabilized by incubation for a short period of time in a solution containing low concentrations of detergents in a hypotonic media. Useful detergents include Nonidet-P 40 (NP40), Triton X-100 (TX-100) or deoxycholate at concentrations of about 0.01 nM to 1.0 mM, generally between about 0.1 μM to about 0.01 mM, particularly about 1 μM . Such systems can be formed from intact cells in culture, including bacterial cells, primary cells, immortalized cell lines, human cells or mixed cell populations.

[0127] A target polypeptide can be obtained from a host cell transformed with and expressing a nucleic acid encoding the target polypeptide. The target nucleic acid can be amplified, for example, by PCR, inserted into an expression vector, and the expression vector introduced into a host cell suitable for expressing the polypeptide encoded by the target nucleic acid. Host cells can be eukaryotic cells, particularly mammalian cells such as human cells, or prokaryotic cells, including, for example, *E. coli*. Eukaryotic and prokaryotic expression vectors are well known in the art and can be obtained from commercial sources. Following expression in the host cell, the target polypeptide can be isolated using methods as disclosed herein. For example, if the target polypeptide is fused to a His-6 peptide, the target polypeptide can be purified by affinity chromatography on a chelated nickel ion column.

AMPLIFICATION OF THE TARGET NUCLEIC ACID SEQUENCE

[0128] At least a portion of a target nucleic acid can be amplified prior to obtaining the target polypeptide encoded by the nucleic acid. PCR, for example, can be performed prior to *in vitro* transcription and translation of a target nucleic acid. Amplification processes include the polymerase chain reaction (Newton and Graham, "PCR" (BIOS Publ. 1994)); nucleic acid sequence based amplification; transcription-based amplification system, self-sustained sequence replication; Q-beta replicase based amplification; ligation amplification reaction; ligase chain reaction (Wiedmann et al., PCR Meth. Appl., 3:57-64 (1994); Barany, Proc. Natl. Acad. Sci., USA 88, 189-93 (1991)); strand displacement amplification (Walker et al., Nucl. Acids Res. 22:2670-77 (1994)); and variations of these methods, including, for example, reverse transcription PCR (RT-PCR; Higuchi et al., Bio/Technology 11:1026-1030 (1993)), and allele-specific amplification.

[0129] Where a nucleotide sequence of the target nucleic acid is amplified by PCR, well known reaction conditions are used. The minimal components of an amplification reaction include a template DNA molecule; a forward primer and a reverse primer, each of which is capable of hybridizing to the template DNA molecule or a nucleotide sequence linked thereto; each of the four different nucleoside triphosphates or appropriate analogs thereof; an agent for polymerization such as DNA polymerase; and a buffer having the appropriate pH, ionic strength, cofactors, and the like. Generally, about 25 to 30 amplification cycles, each including a denaturation step, an annealing step and an extension step, are performed, but fewer cycles can be sufficient or more cycles can be required depending, for example, on the amount of the template DNA molecules present in the reaction. Examples of PCR reaction conditions are described in U.S. Patent

No. 5,604,099.

[0130] A nucleic acid sequence can be amplified using PCR as described in U.S. Patent No. 5,545,539, which provides an improvement of the basic procedure for amplifying a target nucleotide sequence by including an effective amount of a glycine-based osmolyte in the amplification reaction mixture. The use of a glycine-based osmolyte improves amplification of sequences rich in G and C residues and, therefore, can be useful, for example, to amplify trinucleotide repeat sequences such as those associated with Fragile X syndrome (CGG repeats) and myotonic dystrophy (CTG repeats)

[0131] A primer can be prepared from a naturally occurring nucleic acid, for example, by purification from a restriction digest of the nucleic acid, or can be produced synthetically. A primer is capable of acting as a point of initiation of nucleic acid synthesis when placed under conditions sufficient for synthesis of a primer extension product. Particularly useful primers can hybridize specifically to the target sequence or to sequences adjacent to the target sequence.

[0132] Any specific nucleic acid sequence can be amplified by PCR. It is only necessary that a sufficient number of bases at the ends of the target sequence or in the target sequence be known so as to allow preparation of two oligonucleotide primers that can hybridize to the termini of the sequence to be amplified and its complement, at relative positions along each sequence such that an extension product synthesized from one primer, when it is separated from its template (complement), can serve as a template for extension from the other primer into a nucleic acid of defined length. The greater the knowledge about the bases at both ends of the sequence, the greater can be the specificity of the primers for the target nucleic acid sequence and, therefore, the greater the efficiency of the amplification process. If desired, however, a primer specific for one end of the target nucleic acid can be used and a second primer, based on a known sequence linked to the opposite terminus of the target nucleic acid, can be used for amplification of the complementary strand.

[0133] A primer must be sufficiently long to prime the synthesis of extension products in the presence of the agent for polymerization. The exact length of a primer will depend on many factors, including the temperature at which hybridization and primer extension are to be performed; the composition of the primer; and the method used. Depending on the complexity of the target sequence, a primer generally contains about 9 to about 25 nucleotides, although it can contain more nucleotides. As compared to longer primers, shorter primers generally require lower temperatures to form sufficiently stable hybrid complexes with a template nucleic acid (see Sambrook et al., Molecular Cloning: A laboratory manual (Cold Spring Harbor Laboratory Press 1989).

[0134] Primers as disclosed herein are selected to be substantially complementary to the different strands of each specific sequence to be amplified. As such, the primers can hybridize specifically with their respective complementary strands under defined hybridization conditions. A primer sequence need not reflect the exact sequence of the template. For example, a non-complementary nucleotide fragment can be attached to the 5' end of the primer, with the remainder of the primer sequence being complementary to the template strand. Primers generally should have exact complementarity with a sequence from the target nucleic acid, or complement thereof, so that optimal amplification can be obtained.

[0135] A forward or the reverse primer can contain, if desired, a nucleotide sequence of a promoter, for example, a bacteriophage promoter such as an SP6, T3 or T7 promoter. Amplification of a target nucleic sequence using such a primer produces an amplified target nucleic acid operably linked to the promoter. Such a nucleic acid can be used in an *in vitro* transcription reaction to transcribe the amplified target nucleic acid sequence. Nucleotide sequences of the SP6, T3 and T7 promoter are set forth below:

- SP6 promoter sequences:

5' d(CATACGATTTAGGTGACACTATAG)3' SEQ ID NO: 1;
5' d(ATTAGGTGACACTATAG)3' SEQ ID NO: 2;

- T3 promoter sequence:

5' d(ATTAAACCCTCACTAAAGGGA)3' SEQ ID NO: 3;

and

- T7 promoter sequence:

5' d(TAATACGACTCACTATAGGG)3' SEQ ID NO: 4.

[0136] A primer, which can contain a promoter, also can contain an initiation (ATG) codon, or complement thereof, as appropriate, located downstream of the promoter, such that amplification of the target nucleic acid results in an amplified target sequence containing an ATG codon in frame with the desired reading frame. The reading frame can be the natural reading frame or can be any other reading frame. Where the target polypeptide does not exist naturally, operably linking an initiation codon to the nucleic acid encoding the target polypeptide allows translation of the target

polypeptide in the desired reading frame.

[0137] A forward or reverse primer also can contain a nucleotide sequence, or the complement of a nucleotide sequence (if present in the reverse primer), encoding a second polypeptide. The second polypeptide can be a tag peptide, which interacts specifically with a particular reagent, for example, an antibody. A second polypeptide also can have an unblocked and reactive amino terminus or carboxyl terminus.

[0138] The fusion of a tag peptide to a target polypeptide or other polypeptide of interest allows the detection and isolation of the polypeptide. A target polypeptide encoded by a target nucleic acid fused to a sequence encoding a tag peptide can be isolated from an *in vitro* translation reaction mixture using a reagent that interacts specifically with the tag peptide, then the isolated target polypeptide can be subjected to mass spectrometry, as disclosed herein. It should be recognized that an isolated target polypeptide fused to a tag peptide or other second polypeptide is in a sufficiently purified form to allow mass spectrometric analysis, since the mass of the tag peptide will be known and can be considered in the determination.

[0139] Numerous tag peptides and the nucleic acid sequences encoding such tag peptides, generally contained in a plasmid, are known and are commercially available (e.g., NOVAGEN). Any peptide can be used as a tag, provided a reagent such as an antibody that interacts specifically with the tag peptide is available or can be prepared and identified. Frequently used tag peptides include a myc epitope, which includes a 10 amino acid sequence from c-myc (see Ellison et al., J. Biol. Chem. 266:21150-21157 (1991)); the pFLAG system (International Biotechnologies, Inc.); the pEZZ-protein A system (Pharmacia); a 16 amino acid peptide portion of the *Haemophilus influenza* hemagglutinin protein; a glutathione-S-transferase (GST) protein; and a His-6 peptide. Reagents that interact specifically with a tag peptide also are known, and some are commercially available and include antibodies and various other molecules, depending on the tag, for example, metal ions such as nickel or cobalt ions, which interact specifically with a polyhistidine peptide such as His-6; or glutathione, which can be conjugated to a solid support such as agarose and interacts specifically with GST.

[0140] A second polypeptide also can be designed to serve as a mass modifier of the target polypeptide encoded by the target nucleic acid. Accordingly, a target polypeptide can be mass modified by translating an RNA molecule encoding the target polypeptide operably linked to a mass modifying amino acid sequence, where the mass modifying sequence can be at the amino terminus or the carboxyl terminus of the fusion polypeptide. Modification of the mass of the polypeptide derived from the target nucleic acid is useful, for example, when several peptides are analyzed in a single mass spectrometric analysis, since mass modification can increase resolution of a mass spectrum and allow for analysis of two or more different target polypeptides by multiplexing.

[0141] A mass modification includes modifications such as, but not limited to, addition of a peptide or polypeptide fragment to the target polypeptide. For example, a target polypeptide can be mass modified by translating the target polypeptide to include additional amino acids, such as polyhistidine, polylysine or polyarginine. These modifications serve not only to aid in mass spectrometric analyses, but also can aid in purification, identification, immobilization. The modifications can be added post-translationally or can be included in the nucleic acid that encodes the resulting polypeptide.

[0142] In addition, where a plurality of target polypeptides is to be differentially mass modified, each target polypeptide in the plurality can be mass modified using a different polyhistidine sequence, for example, His-4, His-5, His-6, and so on. The use of such a mass modifying moiety provides the further advantage that the moiety acts as a tag peptide, which can be useful, for example, for isolating the target polypeptide attached thereto.

[0143] An advantage of the above processes is that they permit multiplexing to be performed on a plurality of polypeptides, and, therefore, are useful for determining the amino acid sequences of each of a plurality of polypeptides, particularly a plurality of target polypeptides.

[0144] More than one target nucleic acid can be amplified in the same reaction using several pairs of primers, each pair of which amplifies a different target nucleic acid sequence in a mixture of starting nucleic acids. Amplification can be performed simultaneously, provided the annealing temperature of all the primer pairs is sufficiently similar, or can be performed sequentially, starting with a first pair of primers having the lowest annealing temperature of several pairs of primers, then, after amplifying the first target nucleic acid, adding a second pair of primers having a higher annealing temperature and performing the second amplification at the higher temperature, and so on. Individual reactions with different primer pairs also can be performed, then the reaction products can be pooled. Using such methods provide a means for simultaneously determining the identity of more than one allelic variant of one or more polymorphic regions of one or more genes or genetic lesion.

[0145] A primer, for example, the forward primer, also can contain regulatory sequence elements necessary for translation of an RNA in a prokaryotic or eukaryotic system. In particular, where it is desirable to perform a translation reaction in a prokaryotic translation system, a primer can contain a prokaryotic ribosome binding sequence (Shine-Dalgarno sequence) located downstream of a promoter sequence and about 5 to 10 nucleotides upstream of the initiation codon. A prokaryotic ribosome binding sequence, for example, can have the nucleotide sequence, TAAGGAGG (SEQ ID NO: 5).

[0146] A primer, generally the reverse primer, also can contain a sequence encoding a STOP codon in one or more of the reading frames, to assure proper termination of the target polypeptide. Further, by incorporating into the reverse

primer sequences encoding three STOP codons, one into each of the three possible reading frames, optionally separated by several residues, additional mutations that occur downstream (3') of a mutation that otherwise results in premature termination of a polypeptide can be detected.

[0147] For preparing the primers for the amplification process, the nucleotide sequences of numerous target nucleic acids can be obtained from GenBank, or from relevant journal articles, patents or published patent applications. Oligonucleotide primers can be prepared using any suitable method, including, for example, organic synthesis of a nucleic acid from nucleoside derivatives, and can be performed in solution or on a solid support. The phosphotriester method, for example, has been utilized to prepare gene fragments or short genes. In the phosphotriester method, oligonucleotides are prepared, then joined together to form longer nucleic acids (see Narang et al., Meth. Enzymol. 68:90 (1979); U.S. Patent No. 4,356,270). Primers also can be synthesized as described in U.S. Patent No. 5,547,835; U.S. Patent No. 5,605,798 or U.S. Patent No. 5,622,824.

[0148] Primers for amplification are selected such that the amplification reaction produces a nucleic acid that, upon transcription and translation, can result in a non-naturally occurring polypeptide, for example, a polypeptide encoded by an open reading frame that is not the open reading frame encoding the natural polypeptide. Accordingly, by appropriate primer design, in particular, by including an initiation codon in the desired reading frame and, if present, downstream of a promoter in the primer, a polypeptide produced from a target nucleic acid can be encoded by one of the two non-coding frames of the nucleic acid. Such a method can be used to shift out of frame STOP codons, which prematurely truncate the protein and exclude relevant amino acids, or to make a polypeptide containing an amino acid repeat more soluble.

[0149] A non-naturally occurring target polypeptide also can be encoded by a 5' or 3' non-coding region of an exonic region of a nucleic acid; by an intron; or by a regulatory element such as a promoter sequence that contains, in one of the six frames (3 frames per strand), at least a portion of an open reading frame. In these situations, one primer for amplification of the target nucleic acid contains a promoter and an initiation codon, such that the amplified nucleic acid can be transcribed and translated *in vitro*. Thus, a method for determining the identity of a target polypeptide, as disclosed herein, permits the determination of the identity of a nucleotide sequence located in any region of a chromosome, provided a polypeptide of at least 2 amino acids, generally at least 3 or 4 amino acids, particularly at least 5 amino acids, is encoded by one of the six frames of the polynucleotide.

IMMOBILIZATION OF A POLYPEPTIDE TO A SOLID SUPPORT

[0150] For mass spectrometric analyses, a target polypeptide or other polypeptide of interest can be conjugated and immobilized to a solid support in order to facilitate manipulation of the polypeptide. Such supports are well known to those of skill in the art, and include any matrix used as a solid support for linking proteins. The support is selected to be impervious to the conditions of mass spectrometric analyses. Supports, which can have a flat surface or a surface with structures, include, but are not limited to, beads such as silica gel beads, controlled pore glass beads, magnetic beads, Dynabeads, Wang resin; Merrifield resin, SEPHADEX/SEPHAROSE beads or cellulose beads; capillaries; flat supports such as glass fiber filters, glass surfaces, metal surfaces (including steel, gold silver, aluminum, silicon and copper), plastic materials (including multiwell plates or membranes (formed, for example, of polyethylene, polypropylene, polyamide, polyvinylidene difluoride), wafers, combs, pins or needles (including arrays of pins suitable for combinatorial synthesis or analysis) or beads in an array of pits; wells, particularly nanoliter wells, in flat surfaces, including wafers such as silicon wafers; and wafers with pits, with or without filter bottoms. A solid support is appropriately functionalized for conjugation of the polypeptide and can be of any suitable shape appropriate for the support.

[0151] A solid support, such as a bead, can be functionalized for the immobilization of polypeptides, and the bead can be further associated with a solid support, if desired. Where a bead is to be conjugated to a second solid support, polypeptides can be immobilized on the functionalized support before, during or after the bead is conjugated to the second support.

[0152] A polypeptide of interest can be conjugated directly to a solid support or can be conjugated indirectly through a functional group present either on the support, or a linker attached to the support, or the polypeptide or both. For example, a polypeptide can be immobilized to a solid support due to a hydrophobic, hydrophilic or ionic interaction between the support and the polypeptide. Although such a method can be useful for certain manipulations such as for conditioning of the polypeptide prior to mass spectrometry, such a direct interaction is limited in that the orientation of the polypeptide is not known and can be random based on the position of the interacting amino acids, for example, hydrophobic amino acids, in the polypeptide. Thus, a polypeptide generally is immobilized in a defined orientation by conjugation through a functional group on either the solid support or the polypeptide or both.

[0153] A polypeptide of interest can be modified by adding an appropriate functional group to the carboxyl terminus or amino terminus of the polypeptide, or to an amino acid in the peptide, for example, to a reactive side chain, or to the peptide backbone. It should be recognized, however, that a naturally occurring amino acid normally present in the polypeptide also can contain a functional group suitable for conjugating the polypeptide to the solid support. For example,

a cysteine residue present in the polypeptide can be used to conjugate the polypeptide to a support containing a sulfhydryl group, for example, a support having cysteine residues attached thereto, through a disulfide linkage. Other bonds that can be formed between two amino acids, include, for example, monosulfide bonds between two lanthionine residues, which are non-naturally occurring amino acids that can be incorporated into a polypeptide; a lactam bond formed by a transamidation reaction between the side chains of an acidic amino acid and a basic amino acid, such as between the γ -carboxyl group of Glu (or β -carboxyl group of Asp) and the ϵ -amino group of Lys; or a lactone bond produced, for example, by a crosslink between the hydroxy group of Ser and the γ -carboxyl group of Glu (or β -carboxyl group of Asp). Thus, a solid support can be modified to contain a desired amino acid residue, for example, a Glu residue, and a polypeptide having a Ser residue, particularly a Ser residue at the carboxyl terminus or amino terminus, can be conjugated to the solid support through the formation of a lactone bond. It should be recognized, however, that the support need not be modified to contain the particular amino acid, for example, Glu, where it is desired to form a lactone-like bond with a Ser in the polypeptide, but can be modified, instead, to contain an accessible carboxyl group, thus providing a function corresponding to the γ -carboxyl group of Glu.

[0154] A polypeptide of interest also can be modified to facilitate conjugation to a solid support, for example; by incorporating a chemical or physical moiety at an appropriate position in the polypeptide, generally the C-terminus or N-terminus. The artisan will recognize, however, that such a modification, for example, the incorporation of a biotin moiety, can affect the ability of a particular reagent to interact specifically with the polypeptide and, accordingly, will consider this factor, if relevant, in selecting how best to modify as polypeptide of interest.

[0155] In one aspect of the processes provided herein, a polypeptide of interest can be covalently conjugated to a solid support and the immobilized polypeptide can be used to capture a target polypeptide, which binds to the immobilized polypeptide. The target polypeptide then can be released from immobilized polypeptide by ionization or volatilization for mass spectrometry, whereas the covalently conjugated polypeptide remains bound to the support.

[0156] Accordingly, a method to determine the identity of polypeptides that interact specifically with a polypeptide of interest is provided. For example, such a process can be used to determine the identity of target polypeptides obtained from one or more biological samples that interact specifically with the immobilized polypeptide of interest. Such a process also can be used, for example, to determine the identity of binding proteins such as antibodies that bind to the immobilized polypeptide antigen of interest, or receptors that bind to an immobilized polypeptide ligand of interest, or the like. Such a process can be useful, for example, for screening a combinatorial library of modified target polypeptides such as modified antibodies, antigens, receptors, hormones, or other polypeptides to determine the identity of those target polypeptides that interact specifically with the immobilized polypeptide.

[0157] In one aspect of the processes provided herein, a polypeptide of interest can be covalently conjugated to a solid support and the immobilized polypeptide can be used to capture a target polypeptide, which binds to the immobilized polypeptide. The target polypeptide then can be released from immobilized polypeptide by ionization or volatilization for mass spectrometry, whereas the covalently conjugated polypeptide remains bound to the support.

[0158] Accordingly, a process is provided to determine the identity of polypeptides that interact specifically with a polypeptide of interest. For example, such a process can be used to determine the identity of target polypeptides obtained from one or more biological samples that interact specifically with the immobilized polypeptide of interest. Such a process also can be used, for example, to determine the identity of binding proteins such as antibodies that bind to the immobilized polypeptide antigen of interest, or receptors that bind to an immobilized polypeptide ligand of interest, or the like. Such a process can be useful, for example, for screening a combinatorial library of modified target polypeptides such as modified antibodies, antigens, receptors, hormones, or other polypeptides to determine the identity of those target polypeptides that interact specifically with the immobilized polypeptide.

[0159] A polypeptide of interest can be conjugated to a solid support, which can be selected based on advantages that can be provided. Conjugation of a polypeptide to a support, for example, provides the advantage that a support has a relatively large surface area for immobilization of polypeptides. A support, such as a bead, can have any three dimensional structure, including a surface to which a polypeptide, functional group, or other molecule can be attached. If desired, a support, such as a bead, can have the additional characteristic that it can be conjugated further to a different solid support, for example, to the walls of a capillary tube. A support useful for the disclosed processes generally has a size in the range of about 1 to about 100 μm in diameter; can be made of any insoluble or solid material, as disclosed above; and can be a swellable bead, for example, a polymeric bead such as Wang resin, or a non-swellable bead such as a controlled pore glass.

[0160] A solid surface also can be modified to facilitate conjugation of a polypeptide of interest. A thiol-reactive functionality is particularly useful for conjugating a polypeptide to a solid support. A thiol-reactive functionality is a chemical group that can rapidly react with a nucleophilic thiol moiety to produce a covalent bond, for example, a disulfide bond or a thioether bond. In general, thiol groups are good nucleophiles and, therefore, thiol-reactive functionalities generally are reactive electrophiles. A variety of thiol-reactive functionalities are known in the art, including, for example, haloacetyls such as iodoacetyl; diazoketones; epoxy ketones, α - and β -unsaturated carbonyls such as α -enones and β -enones; and other reactive Michael acceptors such as maleimide; acid halides; benzyl halides; and the like. A free thiol group of

a disulfide, for example, can react with a free thiol group by disulfide bond formation, including by disulfide exchange. Reaction of a thiol group can be temporarily prevented by blocking with an appropriate protecting group, as is conventional in the art (see Greene and Wuts "Protective Groups in Organic Synthesis" 2nd ed. (John Wiley & Sons 1991)).

[0161] Reducing agents that are useful for reducing a polypeptide containing a disulfide bond include tris-(2-carboxyethyl)phosphine (TCEP), which generally is used in a concentration of about 1 to 100 mM, usually about 10 mM, and is reacted at a pH of about 3 to 6, usually about pH 4.5, a temperature of about 20 to 45°C, usually about 37°C, for about 1 to 10 hours, usually about 5 hours; dithiothreitol, which generally is used in a concentration of about 25 to 100 mM, and is reacted at a pH of about 6 to 10, usually about pH 8, a temperature of about 25 to 45°C, usually about 37°C, for about 1 to 10 hours, usually about 5 hours. TCE provides an advantage in that it is reactive at a low pH, which effectively protonates thiols, thus suppressing nucleophilic reactions of thiols and resulting in fewer side reactions than with other disulfide reducing agents.

[0162] A thiol-reactive functionality such as 3-mercaptopropyltriethoxysilane can be used to functionalize a silicon surface with thiol groups. The amino functionalized silicon surface then can be reacted with a heterobifunctional reagent such as N-succinimidyl (4-iodoacetyl) aminobenzoate (SIAB) (Pierce; Rockford IL). If desired, the thiol groups can be blocked with a photocleavable protecting group, which then can be selectively cleaved, for example, by photolithography, to provide portions of a surface activated for immobilization of a polypeptide of interest. Photocleavable protecting groups are known in the art (see, for example, published International PCT application No. WO 92/10092; McCray et al., Ann. Rev. Biophys. Biophys. Chem. 18:239-270 (1989)) and can be selectively deblocked by irradiation of selected areas of the surface using, for example, a photolithography mask.

LINKERS

[0163] As noted herein, the polypeptide can be linked either directly to the support or via a linking moiety or moieties. Any linkers known to those of skill in the art to be suitable for linking peptides or amino acids to supports, either directly or via a spacer, may be used. Linkers, include, Rink amide linkers (see, e.g., Rink (1976) Tetrahedron Letters 28:3787), trityl chloride linkers (see, e.g., Leznoff (1978) Acc. Chem. Res. 11:327), Merrifield linkers (see, e.g., Bodansky et al. (1976) Peptide Synthesis, Academic Press, 2nd edition, New York). For example, trityl linkers are known (see, e.g., U.S. Patent No. 5,410,068 and U.S. Patent No. 5,612,474). Amino trityl linkers (see, Figure 3) are also known (see, e.g., U.S. Patent No. 5,198,531). Linkers that are suitable for chemically linking peptides to supports, include disulfide bonds, thioether bonds, hindered disulfide bonds, and covalent bonds between free reactive groups, such as amine and thiol groups. These bonds can be produced using heterobifunctional reagents to produce reactive thiol groups on one or both of the polypeptides and then reacting the thiol groups on one polypeptide with reactive thiol groups or amine groups on the other. Other linkers include, acid cleavable linkers, such as bismaleimideoxy propane, acid labile-transferrin conjugates and adipic acid dihydrazide, that would be cleaved in more acidic intracellular compartments; photocleavable cross linkers that are cleaved by visible or UV light, RNA linkers that are cleavable by ribozymes and other RNA enzymes, and linkers, such as the various domains, such as C_H1, C_H2, and C_H3, from the constant region of human IgG₁ (see, Batra et al. (1993) Molecular Immunol. 30:379-386).

[0164] Any linker known to one skilled in the art for immobilizing a polypeptide to a solid support can be used in a process as disclosed herein. Combinations of any linkers are also contemplated herein. For example, a linker that is cleavable under mass spectrometric conditions, such as a silyl linkage or photocleavable linkage, can be combined with a linker, such as an avidin biotin linkage, that is not cleaved under these conditions, but may be cleaved under other conditions.

[0165] A polypeptide of interest can be attached directly to a support or via a linker. For example, the polypeptide can be conjugated to a support, such as a bead, through means of a variable spacer. In addition, the conjugation can be directly cleavable, for example, through a photocleavable linkage such as a streptavidin or avidin to biotin interaction, which can be cleaved by a laser as occurs for mass spectrometry, or indirectly through a photocleavable linker (see U.S. Patent No. 5,643,722) or an acid labile linker, heat sensitive linker, enzymatically cleavable linker or other such linker.

[0166] A linker can provide a reversible linkage such that it is cleaved under the conditions of mass spectrometry. Such a linker can be, for example, a photocleavable bond such as a charge transfer complex or a labile bond formed between relatively stable organic radicals. A linker (L) on a polypeptide can form a linkage, which generally is a temporary linkage, with a second functional group (L') on the solid support. Furthermore, where the polypeptide of interest has a net negative charge, or is conditioned to have such a charge, the linkage can be formed with L' being, for example, a quaternary ammonium group. In this case, the surface of the solid support carries a negative charge that repels the negatively charged polypeptide, thereby facilitating desorption of the polypeptide for mass spectrometric analysis. Desorption can occur due to the heat created by the laser pulse or, where L' is a chromophore, by specific absorption of laser energy that is in resonance with the chromophore.

[0167] A linkage (L-L') can be, for example, a disulfide bond, which is chemically cleavable by mercaptoethanol or dithioerythrol; a biotin/streptavidin linkage, which can be photocleavable; a heterobifunctional derivative of a trityl ether

group, which can be cleaved by exposure to acidic conditions or under conditions of mass spectrometry (Köster et al., "A Versatile Acid-Labile Linker for Modification of Synthetic Biomolecules," *Tetrahedron Lett.* 31:7095 (1990)); a levulinyl-mediated linkage, which can be cleaved under almost neutral conditions with a hydrazinium/acetate buffer; an arginine-arginine or a lysine-lysine bond, either of which can be cleaved by an endopeptidase such as trypsin; a pyrophosphate bond, which can be cleaved by a pyrophosphatase; or a ribonucleotide bond, which can be cleaved using a ribonuclease or by exposure to alkali condition.

[0168] The functionalities, Land L', can also form a charge transfer complex, thereby forming a temporary L-L' linkage. Since the "charge-transfer band" can be determined by UV/vis spectrometry (see Foster, "Organic Charge Transfer Complexes" (Academic Press 1969)), the laser energy can be tuned to the corresponding energy of the charge-transfer wavelength and specific desorption from the solid support can be initiated. It will be recognized that several combinations of L and L' can serve this purpose and that the donor functionality can be on the solid support or can be coupled to the polypeptide to be detected or vice versa.

[0169] A reversible L-L' linkage also can be generated by homolytically forming relatively stable radicals. Under the influence of the laser pulse, desorption, as well as ionization, can take place at the radical position. Various organic radicals can be selected such that, in relation to the dissociation energy needed to homolytically cleave the bond between the radicals, a corresponding laser wavelength can be selected (see Wentrup, "Reactive Molecules" (John Wiley & Sons 1984)).

[0170] Other linkers include are those that can be incorporated into fusion proteins and expressed in a host cell. Such linkers may be selected amino acids, enzyme substrates, or any suitable peptide. The linker may be made, for example, by appropriate selection of primers when isolating the nucleic acid. Alternatively, they may be added by post translational modification of the protein of interest.

[0171] In particular, selectively cleavable linkers, including photocleavable linkers, acid cleavable linkers, acid-labile linkers, and heat sensitive linkers are useful. Acid cleavable linkers include, for example, bismaleimideoxy propane, adipic acid dihydrazide linkers (see Fattom et al., *Infect. Immun.* 60:584-589 (1992)), and acid labile transferrin conjugates that contain a sufficient portion of transferrin to permit entry into the intracellular transferrin cycling pathway (see Welhöner et al., *J. Biol. Chem.* 266:4309-4314 (1991)).

[0172] FIGURE 2 shows a preferred embodiment of a method of orthogonal capture, cleavage and MALDI analysis of a peptide. This embodiment demonstrates capture through the amino-terminus of the peptide. As shown, the peptide is captured onto a surface of a support through the use of a diisopropylsilyl diether group. Other silyl diether groups, including, but not limited to, dialkylsilyl, diarylsilyl and alkylarylsilyl, may also be used. Reaction of a hydroxylated support surface with diisopropylsilyl dichloride and a hydroxyester provides the starting surface-bound diisopropylsilyl diether ester.

[0173] With reference to the FIGURE, R³ is any attachment moiety, resulting from a support that has been derivatized for linkage, with a derivatizing group that has a hydroxyl group available for reaction. R³ also can be a linkage, such as biotin-streptavidin or biotin-avidin. R³ includes groups such as polyethylene glycol (PEG), an alkylene or arylene group.

[0174] The hydroxylated support surface may be prepared by methods that are well-known to those of skill in the art. For example, N-succinimidyl(4-iodoacetyl) aminobenzoate (SIAB). Other agents as linkers (R³) include, but are not limited to, dimaleimide, dithio-bis-nitrobenzoic acid (DTNB), N-succinimidyl-S-acetyl-thioacetate (SATA), N-succinimidyl-3-(2-pyridyldithiol propionate (SPDP), succinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate (SMCC) and 6-hydrazinonicotimide (HYNIC) may also be used in the novel process. For further examples of cross-linking reagents, see, e.g., Wong "Chemistry of Protein Conjugation and Cross-Linking," CRC Press (1991), and Hermanson, "Bioconjugate. Techniques" Academic Press (1995). Hydroxyesters that may be used include, but are not limited to, hydroxyacetate (glycolate), α -, β -, γ -, ..., ω -hydroxyalkanoates, ω -hydroxy(polyethyleneglycol)COOH, hydroxybenzoates, hydroxyarylalkanoates and hydroxyalkylbenzoates. Thus, with reference to FIGURE 2, R⁴ may be any divalent group that is 2 or more bonds in length, such as (CH₂)_n, where n is 2 or more, and polyethylene glycol. The derivatized support is then reacted with the desired peptide to capture the peptide on the support with loss of R¹OH. The peptide may be reacted directly with the ester group in embodiments where COOR¹ is an active ester group. In these preferred embodiments, R¹ is selected from groups such as, but not limited to, N-succinimidyl, sodium 3-sulfo-N-succinimidyl and 4-nitrophenyl. In other embodiments, the ester is saponified, e.g., with hydroxide, to provide the corresponding acid. This acid is then coupled with the amino-terminus of the peptide under standard peptide coupling conditions (e.g., 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) and N-hydroxysuccinimide (NHS)). The captured peptide is then truncated (fragmented) by reaction with an enzyme or reagent specific for a given amide bond of the peptide. Cleavage of the truncated peptide, containing an N-terminal fragment of the original peptide, from the support is then accomplished by reaction with mild acid. Acids suitable for this cleavage include, but are not limited to, acetic acid, trifluoroacetic acid, paratoluenesulfonic acid and mineral acids. A preferred acid is 3-hydroxypicolinic acid, which is also a suitable matrix for the subsequent MALDI analysis.

[0175] FIGURE 3 illustrates other preferred linkers and capture strategies for MALDI analysis of peptides. As shown, the peptide may be captured through the carboxy terminus by employing an amino-derivatized support. The starting


amino-derivatized support may be prepared by reacting a hydroxylated support surface with diisopropylsilyl dichloride and an aminoalcohol. Aminoalcohols that may be used include, but are not limited to, α -, β -, γ -, ..., ω -aminoalkanols, ω -hydroxy(polyethyleneglycol) NH_2 , hydroxyanilines, hydroxyarylalkylamines and hydroxyalkylanilines. Thus, with reference to FIGURE 3, R^4 may be any divalent group that is 2 or more bonds in length. Capture of the peptide by the amino-derivatized support is achieved by dehydrative coupling of the peptide with the amino group. Such peptide coupling conditions are well-known to those of skill in the art. Illustrated is one set of conditions for capture of the peptide (i.e., 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) and N-hydroxysuccinimide (NHS)). The captured peptide may then be truncated, cleaved from the support, and analyzed as shown in FIGURE 2.

[0176] Also illustrated in FIGURE 3 are other linkers useful in capturing peptides on supports for MALDI analysis. For example, trityl-containing linkers, functionalized with either ester or amino moieties, may be used to capture peptides at the amino or carboxy terminus, respectively. Other linkers known to those of skill in art, e.g., photocleavable linkers, are also available for use in capturing the peptides on the support surface.

Photocleavable Linkers


[0177] Photocleavable linkers are provided. The linkers contain o-nitrobenzyl moieties and phosphate linkages, which allow for complete photolytic cleavage of the conjugates within minutes upon UV irradiation. The UV wavelengths used are selected so that the irradiation will not damage the polypeptides and generally are about 350 to 380 nm, usually about 365 nm.

[0178] A photocleavable linker can have the general structure of formula I:



where R^{20} is ω -(4,4'-dimethoxytrityloxy)alkyl or ω -hydroxyalkyl; R^{21} is selected from hydrogen, alkyl, aryl, alkoxycarbonyl, aryloxycarbonyl and carboxy; R^{22} is hydrogen or (dialkylamino)(ω -cyanoalkoxy)P-; t is 0-3; and R^{50} is alkyl, alkoxy, aryl or aryloxy.

[0179] A photocleavable linker also can have the formula II:




where R^{20} is ω -(4,4'-dimethoxytrityloxy)alkyl, ω -hydroxyalkyl or alkyl; R^{21} is selected from hydrogen, alkyl, aryl, alkoxycarbonyl, aryloxycarbonyl and carboxy; R^{22} is hydrogen or (dialkylamino)(ω -cyanoalkoxy)P-; and X^{20} is hydrogen, alkyl or OR^{20} .

[0180] In a particular photocleavable linker, R^{20} is 3-(4,4'-dimethoxytrityloxy)propyl, 3-hydroxypropyl or methyl; R^{21} is selected from hydrogen, methyl and carboxy; R^{22} is hydrogen or (diisopropylamino)(2-cyanoethoxy)P-; and X^{20} is hydrogen, methyl or OR^{20} . In another photocleavable, R^{20} is 3-(4,4'-dimethoxytrityloxy)propyl; R^{21} is methyl; R^{22} is

(diisopropylamino) (2-cyanoethoxy)P-; and X²⁰ is hydrogen. In still another photocleavable linker, R²⁰ is methyl; R²¹ is methyl; R²² is (diisopropylamino) (2-cyanoethoxy)P-; and X²⁰ is 3-(4,4'-dimethoxytrityloxy)propoxy.

[0181] A photocleavable linker also can have the general formula of formula III:



where R²³ is hydrogen or (dialkylamino)(ω -cyanoalkoxy)P-; and R²⁴ is selected from ω -hydroxyalkoxy, ω -(4,4'-dimethoxytrityloxy)alkoxy, ω -hydroxyalkyl and ω -(4,4'-dimethoxytrityloxy)alkyl, and is unsubstituted or substituted on the alkyl or alkoxy chain with one or more alkyl groups; r and s are each independently 0-4; and R⁵⁰ is alkyl, alkoxy, aryl or aryloxy.

[0182] In particular photocleavable linkers, R²⁴ is ω -hydroxyalkyl or ω -(4,4'-dimethoxytrityloxy)alkyl, and is substituted on the alkyl chain with a methyl group. In another photocleavable linker, R²³ is hydrogen or (diisopropylamino)(2-cyanoethoxy)P-; and R²⁴ is selected from 3-hydroxypropoxy, 3-(4,4'-dimethoxytrityloxy)propoxy, 4-hydroxybutyl, 3-hydroxy-1-propyl, 1-hydroxy-2-propyl, 3-hydroxy-2-methyl-1-propyl, 2-hydroxyethyl, hydroxymethyl, 4-(4,4'-dimethoxytrityloxy)butyl, 3-(4,4'-dimethoxytrityloxy)-1-propyl, 2-(4,4'-dimethoxytrityloxy) ethyl, 1-(4,4'-dimethoxytrityloxy)-2-propyl, 3-(4,4'-dimethoxytrityloxy)-2-methyl-1-propyl and 4,4'-dimethoxytrityloxymethyl. In still another photocleavable linker, R²³ is (diisopropylamino) (2-cyanoethoxy)P-; r and s are O; and R²⁴ is selected from 3-(4,4'-dimethoxytrityloxy)propoxy; 4-(4,4'-dimethoxytrityloxy)butyl, 3-(4,4'-dimethoxytrityloxy) propyl, 2-(4,4'-dimethoxytrityloxy)ethyl, 1-(4,4'-dimethoxytrityloxy)-2-propyl, 3-(4,4'-dimethoxytrityloxy)-2-methyl-1-propyl and 4,4'-dimethoxytrityloxymethyl. R²⁴ is most preferably 3-(4,4'-dimethoxytrityloxy)propoxy.

Preparation of the photocleavable linkers


Preparation of photocleavable linkers of formulae I or II

[0183] Photocleavable linkers of formulae I or II can be prepared by the methods described below, by minor modification of the methods by choosing the appropriate starting materials or by any other methods known to those of skill in the art. Detailed procedures for the synthesis of photocleavable linkers of formula II are provided in Examples 2 and 3.

[0184] In the photocleavable linkers of formula II, where X²⁰ is hydrogen, the linkers can be prepared in the following manner. Alkylation of 5-hydroxy-2-nitrobenzaldehyde with an ω -hydroxyalkyl halide, for example, 3-hydroxypropyl bromide, followed by protection of the resulting alcohol, for example, as a silyl ether, provides a 5-(ω -silyloxyalkoxy)-2-nitrobenzaldehyde. Addition of an organometallic to the aldehyde affords a benzylic alcohol. Organometallics that can be used include trialkylaluminums (for linkers where R²¹ is alkyl) such as trimethylaluminum; borohydrides (for linkers where R²¹ is hydrogen) such as sodium borohydride; or metal cyanides (for linkers where R²¹ is carboxy or alkoxycarbonyl) such as potassium cyanide. In the case of the metal cyanides, the product of the reaction, a cyanohydrin, is hydrolyzed under either acidic or basic conditions in the presence of either water or an alcohol to afford the compounds of interest.

[0185] The silyl group of the side chain of the resulting benzylic alcohols can be exchanged for a 4,4'-dimethoxytrityl group by desilylation using, for example, tetrabutylammonium fluoride, to give the corresponding alcohol, followed by reaction with 4,4'-dimethoxytrityl chloride. Reaction, for example, with 2-cyanoethyl diisopropylchlorophosphoramidite affords the linkers where R²² is (dialkylamino) (ω -cyanoalkoxy)P-.

[0186] A specific example of a synthesis of a photocleavable linker of formula II is shown in the following scheme, which also demonstrates use of the linker in oligonucleotide synthesis. This scheme is intended to be illustrative only and in no way limits the scope of the methods herein. Experimental details of these synthetic transformations are provided in the Examples.





[0187] Synthesis of the linkers of formula II, where X²⁰ is OR²⁰, 3,4-dihydroxyacetophenone is protected selectively at the 4-hydroxyl by reaction, for example, with potassium carbonate and a silyl chloride. Benzoate esters, propiophenones, butyrophenones, and the like can be used in place of the acetophenone. The resulting 4-silyloxy-3-hydroxyacetophenone then is alkylated at the 3-hydroxyl with an alkyl halide (for linkers where R²⁰ is alkyl) and desilylated, for example, with tetrabutylammonium fluoride to afford a 3-alkoxy-4-hydroxyacetophenone. This compound then is alkylated at the 4-hydroxyl by reaction with an ω-hydroxyalkyl halide, for example, 3-hydroxypropyl bromide, to give a 4-(ω-hydroxyalkoxy)-3-alkoxy acetophenone. The side chain alcohol is then protected as an ester, for example, an acetate. This compound is then nitrated at the 5-position, for example, with concentrated nitric acid to provide the corresponding 2-nitroacetophenones. Saponification of the side chain ester, for example, with potassium carbonate, and reduction of the ketone, for example, with sodium borohydride, in either order gives a 2-nitro-4-(ω-hydroxyalkoxy)-

5-alkoxybenzylic alcohol.

[0188] Selective protection of the side chain alcohol as the corresponding 4,4'-dimethoxytrityl ether is then accomplished by reaction with 4,4'-dimethoxytrityl chloride. Further reaction, for example, with 2-cyanoethyl diisopropylchlorophosphoramidite affords the linkers where R²² is (dialkylamino) (ω-cyanoalkoxy)P-.

[0189] A specific example of the synthesis of a photocleavable linker of formula II is shown the following scheme. This scheme is intended to be illustrative only and in no way limit the scope of the methods herein.





Preparation of photocleavable linkers of formula III

[0190] Photocleavable linkers of formula III can be prepared by the methods disclosed herein, by minor modification of the methods by choosing appropriate starting materials, or by other methods known to those of skill in the art.

[0191] In general, photocleavable linkers of formula III are prepared from ω -hydroxyalkyl- or alkoxyaryl compounds, in particular ω -hydroxy-alkyl or alkoxy-benzenes. These compounds are commercially available, or may be prepared from an ω -hydroxyalkyl halide, for example, 3-hydroxypropyl bromide, and either phenyllithium (for the ω -hydroxyalkyl-benzenes) or phenol (for the ω -hydroxyalkoxybenzenes). Acylation of the ω -hydroxyl group, for example, as an acetate ester, followed by Friedel-Crafts acylation of the aromatic ring with 2-nitrobenzoyl chloride provides a 4-(ω -acetoxy-alkyl or alkoxy)-2-nitro benzophenone. Reduction of the ketone, for example, with sodium borohydride, and saponification of the side chain ester are performed in either order to afford a 2-nitrophenyl-4-(hydroxy-alkyl or alkoxy)phenylmethanol. Protection of the terminal hydroxyl group as the corresponding 4,4'-dimethoxytrityl ether is achieved by reaction with 4,4'-dimethoxytrityl chloride. The benzylic hydroxyl group is then reacted, for example, with 2-cyanoethyl diisopropylchlorophosphoramidite to afford linkers of formula II where R²³ is (dialkylamino) (ω -cyanoalkoxy)P-.

[0192] Other photocleavable linkers of formula III can be prepared by substituting 2-phenyl-1-propanol or 2-phenyl-

methyl-1-propanol for the ω -hydroxy-alkyl or alkoxy-benzenes in the above synthesis. These compounds are commercially available, but also can be prepared by reaction, for example, of phenylmagnesium bromide or benzylmagnesium bromide, with the requisite oxirane (propylene oxide) in the presence of catalytic cuprous ion.

5 Chemically cleavable linkers

[0193] A variety of chemically cleavable linkers also can be used to link a polypeptide to a solid support. Acid-labile linkers are particularly useful chemically cleavable linkers for mass spectrometry, especially for MALDI-TOF, because the acid labile bond is cleaved during conditioning of the target polypeptide upon addition of a 3-HPA matrix solution.

10 The acid labile bond can be introduced as a separate linker group, for example, an acid labile trityl group, or can be incorporated in a synthetic linker by introducing one or more silyl bridges using diisopropylsilyl, thereby forming a diisopropylsilyl linkage between the polypeptide and the solid support. The diisopropylsilyl linkage can be cleaved using mildly acidic conditions such as 1.5% trifluoroacetic acid (TFA) or 3-HPA/1 % TFA MALDI-TOF matrix solution. Methods for the preparation of diisopropylsilyl linkages and analogs thereof are well known in the art (see, for example, Saha et al., J. Ora. Chem. 58:7827-7831 (1993)).

15 **[0194]** As disclosed herein, a polypeptide of interest can be conjugated to a solid support such as a bead. In addition, a first solid support such as a bead also can be conjugated, if desired, to a second solid support, which can be a second bead or other support, by any suitable means, including those disclosed herein for conjugation of a polypeptide to a support. Accordingly, any of the conjugation methods and means disclosed herein with reference to conjugation of a polypeptide to a solid support also can be applied for conjugation of a first support to a second support, where the first and second solid support can be the same or different.

20 **[0195]** Appropriate linkers, which can be crosslinking agents, for use for conjugating a polypeptide to a solid support include a variety of agents that can react with a functional group present on a surface of the support, or with the polypeptide, or both. Reagents useful as crosslinking agents include homobifunctional and, in particular, heterobifunctional reagents. 25 Useful bifunctional crosslinking agents include, but are not limited to, N-succinimidyl(4-iodoacetyl) aminobenzoate (SIAB); dimaleimide, dithio-bis-nitrobenzoic acid (DTNB), N-succinimidyl-S-acetyl-thioacetate (SATA), N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP), succinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate (SMCC) and 6-hydrazinonicotimide (HYNIC).

30 **[0196]** A crosslinking agent can be selected to provide a selectively cleavable bond between a polypeptide and the solid support. For example, a photolabile crosslinker such as 3-amino-(2-nitrophenyl)propionic acid (Brown et al., Molec. Divers. 4-12 (1995); Rothschild et al., Nucl. Acids Res. 24:351-66 (1996); U.S. Patent No. 5,643,722) can be employed as a means for cleaving a polypeptide from a solid support. Other crosslinking reagents are well known in the art (see, for example, Wong, "Chemistry of Protein Conjugation and Cross-Linking" (CRC Press 1991); Hermanson, *supra*, 1996).

35 **[0197]** A polypeptide can be immobilized on a solid support such as a bead, through a covalent amide bond formed between a carboxyl group functionalized bead and the amino terminus of the polypeptide or, conversely, through a covalent amide bond formed between an amino group functionalized bead and the carboxyl terminus of the polypeptide.

40 **[0198]** In addition, a bifunctional trityl linker can be attached to the support, for example, to the 4-nitrophenyl active ester on a resin such as a Wang resin, through an amino group or a carboxyl group on the resin via an amino resin. Using a bifunctional trityl approach, the solid support can require treatment with a volatile acid such as formic acid or trifluoroacetic acid to ensure that the polypeptide is cleaved and can be removed. In such a case, the polypeptide can be deposited as a beadless patch at the bottom of a well of a solid support or on the flat surface of a solid support. After addition of a matrix solution, the polypeptide can be desorbed into a mass spectrometer.

45 **[0199]** Hydrophobic trityl linkers also can be exploited as acid-labile linkers by using a volatile acid or an appropriate matrix solution, for example, a matrix solution containing 3-HPA, to cleave an amino linked trityl group from the polypeptide. Acid lability also can be changed. For example, trityl, monomethoxytrityl, dimethoxytrityl or trimethoxytrityl can be changed to the appropriate p-substituted, or more acid-labile tritylamine derivatives, of the polypeptide; i.e. trityl ether and tritylamine bonds to the can be made to the polypeptide. Accordingly, a polypeptide can be removed from a hydrophobic linker, for example, by disrupting the hydrophobic attraction or by cleaving tritylether or tritylamine bonds under acidic conditions, including, if desired, under typical mass spectrometry conditions, where a matrix such as 3-HPA acts as an acid.

50 **[0200]** As disclosed herein, a polypeptide can be conjugated to a solid support, for example, a bead, and the bead, either prior to, during or after conjugation of the polypeptide, can be conjugated to a second solid support, where one or both conjugations result in the formation of an acid-labile bond. For example, use of a trityl linker can provide a covalent or a hydrophobic conjugation, and, regardless of the nature of the conjugation, the trityl group is readily cleaved in acidic conditions. Orthogonally cleavable linkers also can be useful for binding a first solid support, for example, a bead to a second solid support, or for binding a polypeptide of interest to a solid support. Using such linkers, a first solid support, for example, a bead, can be selectively cleaved from a second solid support, without cleaving the polypeptide from the support; the polypeptide then can be cleaved from the bead at a later time. For example, a disulfide linker, which can

be cleaved using a reducing agent such as DTT, can be employed to bind a bead to a second solid support, and an acid cleavable bifunctional trityl group could be used to immobilize a polypeptide to the support. As desired, the linkage of the polypeptide to the solid support can be cleaved first, for example, leaving the linkage between the first and second support intact. Trityl linkers can provide a covalent or hydrophobic conjugation and, regardless of the nature of the conjugation, the trityl group is readily cleaved in acidic conditions.

[0201] A first a solid support such as a bead can be conjugated to a second solid support using the methods, linkages and conjugation means disclosed herein. In addition, a bead, for example, can be bound to a second support through a linking group, which can be selected to have a length and a chemical nature such that high density binding of the beads to the solid support, or high density binding of the polypeptides to the beads, is promoted. Such a linking group can have, for example, "tree-like" structure, thereby providing a multiplicity of functional groups per attachment site on a solid support. Examples of such linking groups include polylysine, polyglutamic acid, penta-erythrole and tris-hydroxy-aminomethane.

[0202] A polypeptide can be conjugated to a solid support, or a first solid support also can be conjugated to a second solid support, through a noncovalent interaction. For example, a magnetic bead made of a ferromagnetic material, which is capable of being magnetized, can be attracted to a magnetic solid support, and can be released from the support by removal of the magnetic field. Alternatively, the solid support can be provided with an ionic or hydrophobic moiety, which can allow the interaction of an ionic or hydrophobic moiety, respectively, with a polypeptide, for example, a polypeptide containing an attached trityl group or with a second solid support having hydrophobic character.

[0203] A solid support also can be provided with a member of a specific binding pair and, therefore, can be conjugated to a polypeptide or a second solid support containing a complementary binding moiety. For example, a bead coated with avidin or with streptavidin can be bound to a polypeptide having a biotin moiety incorporated therein, or to a second solid support coated with biotin or derivative of biotin such as imino-biotin.

[0204] It should be recognized that any of the binding members disclosed herein or otherwise known in the art can be reversed with respect to the examples provided herein. Thus, biotin, for example, can be incorporated into either a polypeptide or a solid support and, conversely, avidin or other biotin binding moiety would be incorporated into the support or the polypeptide, respectively. Other specific binding pairs contemplated for use herein include, but are not limited to, hormones and their receptors, enzymes and their substrates, a nucleotide sequence and its complementary sequence, an antibody and the antigen to which it interacts specifically, and other such pairs known to those skilled in the art.

[0205] Immobilization of one or more polypeptides of interest, particularly target polypeptides, facilitates manipulation of the polypeptides. For example, immobilization of the polypeptides to a solid support facilitates isolation of the polypeptides from a reaction, or transfer of the polypeptides during the performance of a series of reactions. As such, immobilization of the polypeptides can facilitate conditioning the polypeptides or mass modification of the polypeptides prior to performing mass spectrometric analysis.

[0206] Examples of preferred binding pairs or linker/interactions are provided in the Table.

TABLE

LINKER/INTERACTION	EXAMPLES
streptavidin-biotin ^{a,c} /photolabile biotin ^b	biotinylated pin, avidin beads, photolabile biotin polypeptide
hydrophobic ^a	C18-coated pin, tritylated polypeptide
magnetic ^a	electromagnetic pin, streptavidin magnetic beads (e.g., DYNABEADS), biotin polypeptide
acid-labile linker ^b	glass pin, bifunctional trityl-linked DNA
amide bond(s) ^c	silicon wafer, Wang resin, amino-linked polypeptide
disulfide bond ^a	silicon wafer, beads are bound on the flat surface forming arrays or in arrays of nanoliter wells, thiol beads, thiolated polypeptide
photocleavable bond/linker	biotinylated pin/wafer, avidin beads, photolabile biotin polypeptide

(continued)

LINKER/INTERACTION	EXAMPLES
thioether bond ^c	silicon wafer, beads are bound on the flat surface forming arrays or in arrays of nanoliter wells, thiolated peptide
^a - these interactions are reversible. ^b - these non-reversible interactions are rapidly cleaved. ^c - unless cleavable-linkers are incorporated at some point in the scheme, only the complement of the solid-bound DNA can be analyzed in these schemes.	

CONDITIONING A POLYPEPTIDE

[0207] Conditioning of a polypeptide prior to mass spectrometry can increase the resolution of a mass spectrum of the polypeptide, thereby facilitating determining the identity of a target polypeptide. A polypeptide can be conditioned, for example, by treating the polypeptide with a cation exchange material or an anion exchange material, which can reduce the charge heterogeneity of the polypeptide, thereby reducing or eliminating peak broadening. In addition, contacting a polypeptide with an alkylating agent such as alkyl iodide, iodoacetamide, iodoethanol, or 2,3-epoxy-1-propanol, for example, can prevent the formation of disulfide bonds in the polypeptide, thereby increasing resolution of a mass spectrum of the polypeptide. Likewise, charged amino acid side chains can be converted to uncharged derivatives by contacting the polypeptides with trialkylsilyl chlorides, thus reducing charge heterogeneity and increasing resolution of the mass spectrum.

[0208] There are also means of improving resolution, particularly for shorter peptides, by incorporating modified amino acids that are more basic than the corresponding unmodified residues. Such modification in general increases the stability of the polypeptide during mass spectrometric analysis. Also, cation exchange chromatography, as well as general washing and purification procedures which remove proteins and other reaction mixture components away from the target polypeptide, can be used to clean up the peptide after *in vitro* translation and thereby increase the resolution of the spectrum resulting from mass spectrometric analysis of the target polypeptide.

[0209] Conditioning also can involve incorporating modified amino acids into the polypeptide, for example, mass modified amino acids, which can increase resolution of a mass spectrum. For example, the incorporation of a mass modified leucine residue in a polypeptide of interest can be useful for increasing the resolution (e.g., by increasing the mass difference) of a leucine residue from an isoleucine residue, thereby facilitating determination of an amino acid sequence of the polypeptide. A modified amino acid also can be an amino acid containing a particular blocking group, such as those groups used in chemical methods of amino acid synthesis. For example, the incorporation of a glutamic acid residue having a blocking group attached to the side chain carboxyl group can mass modify the glutamic acid residue and, provides the additional advantage of removing a charged group from the polypeptide, thereby further increasing resolution of a mass spectrum of a polypeptide containing the blocked amino acid.

USE OF A PIN TOOL TO IMMOBILIZE A POLYPEPTIDE

[0210] The immobilization of a polypeptide of interest to a solid support using a pin tool can be particularly advantageous. Pin tools include those disclosed herein or otherwise known in the art (see, e.g., copending U.S. application Serial Nos. 08/786,988 and 08/787,639, and International PCT application No. WO 98/20166).

[0211] A pin tool in an array, for example, a 4 X 4 array, can be applied to wells containing polypeptides of interest. Where the pin tool has a functional group attached to each pin tip, or a solid support, for example, functionalized beads or paramagnetic beads, are attached to each pin, the polypeptides in a well can be captured (≥ 1 pmol capacity). During the capture step, the pins can be kept in motion (vertical, 1-2 mm travel) to increase the efficiency of the capture. Where a reaction such as an *in vitro* transcription is being performed in the wells, movement of the pins can increase efficiency of the reaction.

[0212] Polypeptides of interest, particularly target polypeptides, are immobilized due to contact with the pin tool. Further immobilization can result by applying an electrical field to the pin tool. When a voltage is applied to the pin tool, the polypeptides are attracted to the anode or the cathode, depending on their net charge. Such a system also can be useful for isolating the polypeptides, since uncharged molecules remain in solution and molecules having a charge opposite to the net charge of the polypeptides are attracted to the opposite pole (anode or cathode). For more specificity, the pin tool (with or without voltage) can be modified to have conjugated thereto a reagent specific for the polypeptide of interest, such that only the polypeptides of interest are bound by the pins. For example, the pins can have nickel ions attached, such that only polypeptides containing a polyhistidine sequence are bound. Similarly, the pins can have antibodies specific for a target polypeptide attached thereto, or to beads that, in turn, are attached to the pins, such that only the

target polypeptides, which contain the epitope recognized by the antibody, are bound by the pins.

[0213] Different pin conformations include, for example, a solid pin configuration, or pins with a channel or with a hole through the center, which can accommodate an optic fiber for mass spectrometer detection. The pin can have a flat tip or any of a number of configurations, including nanowell, concave, convex, truncated conic or truncated pyramidal, for example, a size 4 to 800 μm across x 100 μm in depth. The individual pins, which can be any size desired, generally are as long as about 10 mm, usually about 5 mm long, and particularly about 1 mm long. The pins and mounting plate can be made of polystyrene, which can be one piece injection molded. Polystyrene is convenient for this use because it can be functionalized readily and can be molded to very high tolerances. The pins in a pin tool apparatus can be collapsible, for example, controlled by a scissor-like mechanism, so that the pins can be brought into closer proximity, reducing the overall size.

[0214] Captured polypeptides can be analyzed by a variety of means including, for example, spectrometric techniques such as UV/VIS, IR, fluorescence, chemiluminescence, NMR spectroscopy, mass spectrometry, or other methods known in the art, or combinations thereof. If conditions preclude direct analysis of captured polypeptides, the polypeptides can be released or transferred from the pins, under conditions such that the advantages of sample concentration are not lost. Accordingly, the polypeptides can be removed from the pins using a minimal volume of eluent, and without any loss of sample. Where the polypeptides are bound to the beads attached to the pins, the beads containing the polypeptides can be removed from the pins and measurements made directly from the beads.

[0215] Prior to determining the identity of a target polypeptide by mass spectrometry, a pin tool having the polypeptide attached thereto can be withdrawn and washed several times, for example, in ammonium citrate to condition the polypeptide prior to addition of matrix. The pins then can be dipped into matrix solution, with the concentration of matrix adjusted such that matrix solution adheres only to the very tips of the pins. Alternatively, the pin tool can be inverted and the matrix solution sprayed onto the tip of each pin using a microdrop device. The polypeptides also can be cleaved from the pins, for example, into a nanowell on a chip, prior to addition of matrix. For analysis directly from the pins, a stainless steel "mask" probe can be fitted over the pins, then the mask probe can be installed in the mass spectrometer.

[0216] Two mass spectrometer geometries can be used for accommodating a pin tool apparatus. A first geometry accommodates solid pins. In effect, the laser ablates a layer of material from the surface of the crystals, such that the resultant ions are accelerated and focused through the ion optics. A second geometry accommodates fibre optic pins, in which the laser strikes the samples from behind. In effect, the laser is focused onto the pin tool back plate and into a short optical fibre about 100 μm in diameter and about 7 mm in length to include thickness of the back plate. This geometry requires that the volatilized sample go through the depth of the matrix/bead mix, slowing and cooling down the ions and resulting in a type of delayed extraction, which can increase the resolution of the analysis (see, e.g., Juhasz et al. (1996) Analysis, Anal. Chem. 68:941-946, see also, e.g., U.S. Patent No. 5,777,325, U.S. Patent No. 5,742,049, U.S. Patent No. 5,654,545, U.S. Patent No. 5,641,959, U.S. Patent No. 5,654,545 and U.S. Patent No. 5,760,393 for descriptions of MALDI and delayed extraction protocols).

[0217] The probe through which the pins are fitted also can be of various geometries. For example, a large probe with multiple holes, one for each pin, can be fitted over the pin tool and the entire assembly is translated in the X-Y axes in the mass spectrometer. The probe also can be a fixed probe with a single hole, which is large enough to give an adequate electric field, but small enough to fit between the pins. The pin tool then is translated in all three axes, with each pin being introduced through the hole for sequential analyses. This latter format is more suitable for a higher density pin tool, for example, a pin tool based on a 384 well or higher density microplate format. These two probes are suitable for the two mass spectrometer geometries, as disclosed above.

[0218] Pin tools can be useful for immobilizing polypeptides of interest in spatially addressable manner on an array. Such spatially addressable or pre-addressable arrays are useful in a variety of processes, including, for example, quality control and amino acid sequencing diagnostics. The pin tools described in the copending applications U.S. application Serial Nos. 08/786,988 and 08/787,639 and International PCT application No. WO 98/20166 are serial and parallel dispensing tools that can be employed to generate multi-element arrays of polypeptides on a surface of the solid support. The array surface can be flat, with beads, or geometrically altered to include wells, which can contain beads. A pin tool that allows the parallel development of a sample array is provided. Such a tool is an assembly of vesicle elements, or pins, where each of the pins can include a narrow interior chamber suitable for holding nanoliter volumes of fluid. Each of the pins fits inside a housing that has an interior chamber. The interior housing can be connected to a pressure source that can control the pressure within the interior housing chamber to regulate the flow of fluid through the interior chamber of the pins, thereby allowing for the controlled dispensing of defined volumes of fluid from the vesicles.

[0219] The pin tool also can include a jet assembly, which can include a capillary pin having an interior chamber, and a transducer element mounted to the pin and capable of driving fluid through the interior chamber of the pin to eject fluid from the pin. In this way, the tool can dispense a spot of fluid to a support surface by spraying the fluid from the pin. The transducer also can cause a drop of fluid to extend from the capillary so that fluid can be passed to the array, or other solid support, by contacting the drop to the surface of the array. The pin tool also can form an array of polypeptides by dispensing the polypeptides in a series of steps, while moving the pin to different locations above the array surface to

form the sample array. The pin tool then can pass prepared polypeptide arrays to a plate assembly that disposes the arrays for analysis by mass spectrometry, which generates a set of spectra signal indicative of the composition of the polypeptides under analysis.

[0220] The pin tool can include a housing having a plurality of sides and a bottom portion having formed therein a plurality of apertures, the walls and bottom portion of the housing defining an interior volume; one or more fluid transmitting vesicles, or pins, mounted within the apertures, having a nanovolume sized fluid holding chamber for holding nanovolumes of fluid, the fluid holding chamber being disposed in fluid communication with the interior volume of the housing, and a dispensing element that is in communication with the interior volume of the housing for selectively dispensing nanovolumes of fluid from the nanovolume sized fluid transmitting vesicles when the fluid is loaded with the fluid holding chambers of the vesicles. This allows the dispensing element to dispense nanovolumes of the fluid onto the surface of the support when the apparatus is disposed over and in registration with the support.

[0221] The fluid transmitting vesicle can have an open proximal end and a distal tip portion that extends beyond the housing bottom portion when mounted within the apertures. In this way the open proximal end can dispose the fluid holding chamber in fluid communication with the interior volume when mounted with the apertures. Optionally, the plurality of fluid transmitting vesicles are removably and replaceably mounted within the apertures of the housing, or alternatively can include a glue seal for fixedly mounting the vesicles within the housing.

[0222] The fluid holding chamber also can include a narrow bore, which is dimensionally adapted for being filled with the fluid through capillary action, and can be sized to fill substantially completely with the fluid through capillary action. The plurality of fluid transmitting vesicles includes an array of fluid delivering needles, which can be formed of metal, glass, silica, polymeric material, or any other suitable material, and, thus, as disclosed herein, also can serve as a solid support.

[0223] The housing also can include a top portion, and mechanical biasing elements for mechanically biasing the plurality of fluid transmitting vesicles into sealing contact with the housing bottom portion. In addition, each fluid transmitting vesicle can have a proximal end portion that includes a flange, and further includes a seal element disposed between the flange and an inner surface of the housing bottom portion for forming a seal between the interior volume and an external environment. The biasing elements can be mechanical and can include a plurality of spring elements each of which are coupled at one end to the proximal end of each the plurality of fluid transmitting vesicles, and at another end to an inner surface of the housing top portion. The springs can apply a mechanical biasing force to the vesicle proximal end to form the seal.

[0224] The housing also can include a top portion, and a securing element for securing the housing top portion to the housing bottom portion. The securing element can include a plurality of fastener-receiving apertures formed within one of the top and bottom portions of the housing, and a plurality of fasteners for mounting within the apertures for securing together the housing top and bottom portions.

[0225] The dispensing element can include a pressure source fluidly coupled to the interior volume of the housing for disposing the interior volume at a selected pressure condition. Moreover, where the fluid transmitting vesicles are to be filled through capillary action, the dispensing element can include a pressure controller that can vary the pressure source to dispose the interior volume of the housing at varying pressure conditions. This allows the controller varying element to dispose the interior volume at a selected pressure condition sufficient to offset the capillary action to fill the fluid holding chamber of each vesicle to a predetermined height corresponding to a predetermined fluid amount. Additionally, the controller can include a fluid selection element for selectively discharging a selected nanovolume fluid amount from the chamber of each the vesicle. In addition, a pressure controller that operates under the controller of a computer program operating on a data processing system to provide variable control over the pressure applied to the interior chamber of the housing is provided.

[0226] The fluid transmitting vesicle can have a proximal end that opens onto the interior volume of the housing, and the fluid holding chamber of the vesicles are sized to substantially completely fill with the fluid through capillary action without forming a meniscus at the proximal open end. Optionally, the apparatus can have plural vesicles, where a first portion of the plural vesicles include fluid holding chambers of a first size and a second portion including fluid holding chambers of a second size, whereby plural fluid volumes can be dispensed.

[0227] The tool also can include a fluid selection element that has a pressure source coupled to the housing and in communication with the interior volume for disposing the interior volume at a selected pressure condition, and an adjustment element that couples to the pressure source for varying the pressure within the interior volume of the housing to apply a positive pressure in the fluid chamber of each the fluid transmitting vesicle to vary the amount of fluid dispensed therefrom. The selection element and adjustment element can be computer programs operating on a data processing system that directs the operation of a pressure controller connected to the interior chamber.

[0228] The pin tool apparatus can be used for dispensing a fluid containing a polypeptide of interest, particularly a target polypeptide, into one or more wells of a multi-well device, which can be a solid support. The apparatus can include a housing having a plurality of sides and a bottom portion having formed therein a plurality of apertures, the walls and bottom portion defining an interior volume, a plurality of fluid transmitting vesicles, mounted within the apertures, having

a fluid holding chamber disposed in communication with the interior volume of the housing, and a fluid selection and dispensing means in communication with the interior volume of the housing for variably selecting an amount of the fluid loaded within the fluid holding chambers of the vesicles to be dispensed from a single set of the plurality of fluid transmitting vesicles. Accordingly, the dispensing means dispenses a selected amount of the fluid into the wells of the multi-well device when the apparatus is disposed over and in registration with the device.

[0229] The fluid dispensing apparatus for dispensing fluid containing a polypeptide of interest into one or more wells of a multi-well device can include a housing having a plurality of sides and top and bottom portions, the bottom portion having formed therein a plurality of apertures, the walls and top and bottom portions of the housing defining an interior volume, a plurality of fluid transmitting vesicles, mounted within the apertures, having a fluid holding chamber sized to hold nanovolumes of the fluid, the fluid holding chamber being disposed in fluid communication with the volume of the housing, and mechanical biasing element for mechanically biasing the plurality of fluid transmitting vesicles into sealing contact with the housing bottom portion.

DETERMINING THE MASS OF THE POLYPEPTIDE BY MASS SPECTROMETRY

[0230] The identity of an isolated target polypeptide is determined by mass spectrometry. For mass spectrometry analysis, the target polypeptide can be solubilized in an appropriate solution or reagent system. The selection of a solution or reagent system, for example, an organic or inorganic solvent, will depend on the properties of the target polypeptide and the type of mass spectrometry performed, and is based on methods well known in the art (see, for example, Vorm et al., Anal. Chem. 66:3281 (1994), for MALDI; Valaskovic et al., Anal. Chem. 67:3802 (1995), for ESI). Mass spectrometry of peptides also is described, for example, in International PCT application No. WO 93/24834 to Chait et al. and U.S. Patent No. 5,792,664.

[0231] A solvent is selected so as to considerably reduce or fully exclude the risk that the target polypeptide will be decomposed by the energy introduced for the vaporization process. A reduced risk of target polypeptide decomposition can be achieved, for example, by embedding the sample in a matrix, which can be an organic compound such as a sugar, for example, a pentose or hexose, or a polysaccharide such as cellulose. Such compounds are decomposed thermolytically into CO₂ and H₂O such that no residues are formed that can lead to chemical reactions. The matrix also can be an inorganic compound such as nitrate of ammonium, which is decomposed essentially without leaving any residue. Use of these and other solvents is known to those of skill in the art (see, e.g., U.S. Patent 5,062,935).

[0232] Mass spectrometer formats for use in analyzing a target polypeptide include ionization (I) techniques, such as, but not limited to, matrix assisted laser desorption (MALDI), continuous or pulsed electrospray (ESI) and related methods such as ionspray or thermospray, and massive cluster impact (MCI). Such ion sources can be matched with detection formats, including linear or non-linear reflectron time-of-flight (TOF), single or multiple quadrupole, single or multiple magnetic sector, Fourier transform ion cyclotron resonance (FTICR), ion trap, and combinations thereof such as ion-trap/time-of-flight. For ionization, numerous matrix/wavelength combinations (MALDI) or solvent combinations (ESI) can be employed. Sub-attomole levels of protein have been detected, for example, using ESI mass spectrometry (Valaskovic, et al., Science 273:1199-1202 (1996)) and MALDI mass spectrometry (Li et al., J. Am. Chem. Soc. 118:1662-1663 (1996)).

[0233] Electrospray mass spectrometry has been described by Fenn et al. (J. Phys. Chem. 88:4451-59 (1984); PCT Application No. WO 90/14148) and current applications are summarized in review articles (Smith et al., Anal. Chem. 62: 882-89 (1990); Ardrey, Electrospray Mass Spectrometry, Spectroscopy Europe 4:10-18 (1992)). MALDI-TOF mass spectrometry has been described by Hillenkamp et al. ("Matrix Assisted UV-Laser Desorption/Ionization: A New Approach to Mass Spectrometry of Large Biomolecules, Biological Mass Spectrometry" (Burlingame and McCloskey, eds., Elsevier Science Publ. 1990), pp. 49-60). With ESI, the determination of molecular weights in femtomole amounts of sample is very accurate due to the presence of multiple ion peaks, all of which can be used for mass calculation.

[0234] If desired, a target polypeptide can be conditioned prior to mass spectrometry, as disclosed herein, thus facilitating identification of the polypeptide.

MALDI

[0235] Matrix assisted laser desorption (MALDI) is preferred among the mass spectrometric methods herein. Methods for performing MALDI are well known to those of skill in the art (see, e.g.,). Numerous methods for improving resolution are also known. For example, resolution in MALDI TOF mass spectrometry can be improved by reducing the number of high energy collisions during ion extraction (see, e.g., Juhasz et al. (1996) Analysis, Anal. Chem. 68:941-946, see also, e.g., U.S. Patent No. 5,777,325, U.S. Patent No. 5,742,049, U.S. Patent No. 5,654,545, U.S. Patent No. 5,641,959, U.S. Patent No. 5,654,545, U.S. Patent No. 5,760,393 and U.S. Patent No. 5,760,393 for descriptions of MALDI and delayed extraction protocols).

AMINO ACID SEQUENCING OF TARGET POLYPEPTIDES

[0236] A process of determining the identity of a target polypeptide using mass spectrometry, as disclosed herein, can be performed by determining the amino acid sequence, or a portion thereof, of a target polypeptide. Amino acid sequencing can be performed, for example, from the carboxyl terminus using carboxypeptidase such as carboxypeptidase Y, carboxypeptidase P, carboxypeptidase A, carboxypeptidase G or carboxypeptidase B, or other enzyme that progressively digests a polypeptide from its carboxyl terminus; or from the N-terminus of the target polypeptide by using the Edman degradation method or using an aminopeptidase such as alanine aminopeptidase, leucine aminopeptidase, pyroglutamate peptidase, dipeptidyl peptidase, microsomal peptidase, or other enzyme that progressively digests a polypeptide from its amino terminus. If desired, the target polypeptide first can be cleaved into peptide fragments using an enzyme such as trypsin, chymotrypsin, Asp-N, thrombin or other suitable enzyme. The fragments then can be isolated and subjected to amino acid sequencing by mass spectrometry. If desired, a tag, for example, a tag peptide, can be conjugated to a fragment of a target polypeptide. Such a conjugation can be performed prior to or following cleavage of the target polypeptide.

[0237] Amino acid sequencing of a target polypeptide can be performed either on the free polypeptide or after immobilizing the polypeptide on a solid support. A target polypeptide can be immobilized on a solid support, for example, by linking the polypeptide to the support through its amino terminus or its carboxyl terminus or directly or via a linker or linkers by methods known to those of skill in the art or as described herein, then treating the immobilized polypeptide with an exopeptidase specific for the unbound terminus. For example, where a target polypeptide is linked to a solid support through its amino terminus, the immobilized polypeptide can be treated with a carboxypeptidase, which sequentially degrades the polypeptide from its carboxyl terminus. Alternatively, where the target polypeptide is linked to a solid support through its carboxyl terminus, the polypeptide can be digested from its amino terminus using, for example, Edman's reagent.

[0238] For amino acid sequencing, the target polypeptide is treated with the protease in a time-limited manner, and released amino acids are identified by mass spectrometry. If desired, degradation of a target polypeptide can be performed in a reactor apparatus (see International Publ. No. WO 94121822, published 29 September 1994), in which the polypeptide can be free in solution and the protease can be immobilized, or in which the protease can be free in solution and the polypeptide can be immobilized. At time intervals or as a continuous stream, the reaction mixture containing a released amino acid is transported to a mass spectrometer for analysis. Prior to mass spectrometric analysis, the released amino acids can be transported to a reaction vessel for conditioning, which can be by mass modification. The determination of the amino acid sequence of the target polypeptide, particularly the identification of an allelic variation in the target polypeptide as compared to a corresponding known polypeptide, can be useful, for example, to determine whether the subject from which the target polypeptide was obtained has or is predisposed to a particular disease or condition.

[0239] The target polypeptide is conditioned by mass modified prior to sequencing. It should be recognized, however, that mass modification of a polypeptide prior to chemical or enzymatic degradation, for example, can influence the rate or extent of degradation. Accordingly, the skilled artisan will know that the influence of conditioning and mass modification on polypeptide degradation should be characterized prior to initiating amino acid sequencing.

[0240] A process as disclosed herein is conveniently performed in a multiplexing format, thereby allowing a determination of the identities of a plurality of two or more target polypeptides in a single procedure. For multiplexing, a population of target polypeptides can be synthesized by *in vitro* translation, where each of the target nucleic acids encoding each of the target polypeptides is translated, in a separate reaction, in the presence of one or more mass modifying amino acids. The population of target polypeptides can be encoded, for example, by target nucleic acids representing the different polymorphic regions of a particular gene. Each of the individual reactions can be performed using one or more amino acids that are differentially mass modified particularly using basic residues. Following translation, each target polypeptide is distinguishable by the particular mass modified amino acid.

[0241] A plurality of target polypeptides also can be obtained, for example, from naturally occurring proteins and examined by multiplexing, provided that each of the plurality of target polypeptides is differentially mass modified. For example, where a plurality of target polypeptides are being examined to determine whether a particular polypeptide is an allelic variant containing either a Gly residue or an Ala residue, the Gly and Ala residues in each polypeptide in the plurality can be mass modified with a mass label specific for that polypeptide. Identification of a Gly or Ala residue having a particular mass can be used to determine the particular polypeptide and the nature of the polymorphism.

[0242] Amino acid modifications can be effected during or after *in vitro* translation of the target polypeptide. For example, any amino acid with a functional group on a side chain can be derivatized using methods known to those of skill in the art. For example, N-succinimidy-3(2-pyridyldithio)propionate (SPDP) can be used to introduce sulfhydryl groups on lysine residues, thereby altering the mass of the polypeptide compared to the untreated polypeptide.

EXEMPLARY USES

[0243] Methods for determining the identity of a target polypeptide are disclosed herein. The identity of the target polypeptide allows information to be obtained regarding the DNA sequence encoding the target polypeptide. The target polypeptide can be from a eukaryote such as a vertebrate, particularly a mammal such as a human, or can be from a prokaryote, including a bacterium or a virus. Generally, the target polypeptide can be from any organism, including a plant.

[0244] A target polypeptide can be immobilized to a solid support, thereby facilitating manipulation of the polypeptide prior to mass spectrometry. For example, a target polypeptide can be translated *in vitro*. Such a method of obtaining a target polypeptide conveniently allows attachment of a tag to the polypeptide, for example, by producing a fusion polypeptide of the target polypeptide and a tag peptides such as a polyhistidine tag. The presence of a tag peptide such as a polyhistidine tag provides a means to isolate the target polypeptide, for example, from the *in vitro* translation reaction, by passing the mixture over a nickel chelate column, since nickel ions interact specifically with a polyhistidine sequence. The target polypeptide then can be captured by conjugation to a solid support, thereby immobilizing the target polypeptide. If general, conjugation of the polypeptide to the solid support can be mediated through a linker, which provides desirable characteristics such as being readily cleavable, for example, chemically cleavable, heat cleavable or photocleavable. As shown in Figure 2, for example, the target polypeptide can be immobilized at its amino terminus to a solid support through a diisopropylsilyl linker, which readily is cleavable under acidic conditions such as when exposed to the mass spectrometry matrix solution 3-HPA. For example, the solid support, or a linker conjugated to the support or a group attached to such a linker, can be in the activated carboxy form such as a sulfo-NHS ester, which facilitates conjugation of the polypeptide through its amino terminus. Furthermore, conjugation of a polypeptide to a solid support can be facilitated by engineering the polypeptide to contain, for example, a string of lysine residues, which increases the concentration of amino groups available to react with an activated carboxyl support. Of course, a polypeptide also can be conjugated through its carboxyl terminus using a modified form of the linker shown in Figure 2 (see Figure 3), or can be conjugated using other linkers as disclosed herein or otherwise known in the art. The immobilized target polypeptide then can be manipulated, for example, by proteolytic cleavage using an endopeptidase or a chemical reagent such as cyanogen bromide, by sequential truncation from its free end using an exopeptidase or a chemical reagent such as Edman's reagent, or by conditioning in preparation for mass spectrometric analysis, for example, by cation exchange to improve mass spectrometric analysis. An advantage of performing such manipulations with an immobilized polypeptide is that the reagents and undesirable reaction products can be washed from the remaining immobilized polypeptide, which then can be cleaved from the solid support in a separate reaction or can be subjected to mass spectrometry, particularly MALDI-TOF, under conditions that cleave the polypeptide from the support, for example, exposure of a polypeptide linked to the support through a photocleavable linker to the MALDI laser:

[0245] For purposes of the conjugation reactions, as well as enzymatic reactions, it is assumed that the termini of a target polypeptide are more reactive than the amino acid side groups due, for example, to steric considerations. However, it is recognized that amino acid side groups can be more reactive than the relevant terminus, in which case the artisan would know that the side group should be blocked prior to performing the reaction of interest. Methods for blocking an amino acid side group are well known and blocked amino acid residues are readily available and used, for example, for chemical synthesis of peptides. Similarly, it is recognized that a terminus of interest of the polypeptide can be blocked due, for example, to a post-translational modification, or can be buried within a polypeptide due to secondary or tertiary conformation. Accordingly, the artisan will recognize that a blocked amino terminus of a polypeptide, for example, must be made reactive either by cleaving the amino terminal amino acid or by deblocking the amino acid. In addition, where the terminus of interest is buried within the polypeptide structure, the artisan will know that the polypeptide, in solution, can be heated to about 70 to 100 °C prior to performing a reaction. It is recognized, for example, that when the reaction to be performed is an enzymatic cleavage, the enzymes selected should be stable at elevated temperatures. Such temperature stable enzymes, for example, thermostable peptidases, including carboxypeptidases and aminopeptidases, are obtained from thermophilic organisms and are commercially available. In addition, where it is desirable not to use heat to expose an otherwise buried terminus of a polypeptide, altering the salt conditions can provide a means to expose the terminus. For example, a polypeptide terminus can be exposed using conditions of high ionic strength, in which case an enzyme such as an exopeptidase is selected based on its tolerance to high ionic strength conditions.

[0246] Depending on the target polypeptide to be detected, the disclosed methods allow the diagnosis, for example, of a genetic disease or chromosomal abnormality; a predisposition to or an early indication of a gene influenced disease or condition such as obesity, atherosclerosis, diabetes or cancer; or an infection by a pathogenic organism, including a virus, bacterium, parasite or fungus; or to provide information relating to identity or heredity based, for example, on an analysis of mini-satellites and micro-satellites, or to compatibility based, for example, on HLA phenotyping.

[0247] A process is provided herein for detecting genetic lesions that are characterized by an abnormal number of trinucleotide repeats, which can range from less than 10 to more than 100 additional trinucleotide repeats relative to the number of repeats, if any, in a gene in a non-affected individual. Diseases associated with such genetic lesions include, for example, Huntington's disease, prostate cancer, SCA-1, Fragile X syndrome (Kremer et al., Science 252:1711-14

(1991); Fu et al., Cell 67:1047-58 (1991); Hirst et al., J. Med. Genet. 28:824-29 (1991), myotonic dystrophy type I (Mahadevan et al., Science 255:1253-55 (1992); Brook et al., Cell 68:799-808 (1992)), Kennedy's disease (also termed spinal and bulbar muscular atrophy; La Spada et al., Nature 352:77079 (1991)); Machado-Joseph disease, and dentatorubral and pallidolusian atrophy. The abnormal number of triplet repeats can be located in any region of a gene, including a coding region, a non-coding region of an exon, an intron, or a promoter or other regulatory element. For example, the expanded trinucleotide repeat associated with myotonic dystrophy occurs in the 3' untranslated region (UTR) of the MTPK gene on chromosome 19. In some of these diseases, for example, prostate cancer, the number of trinucleotide repeats is positively correlated with prognosis of the disease such that a higher number of trinucleotide repeats correlates with a poorer prognosis.

[0248] A process for determining the identity of an allelic variant of a polymorphic region of a gene, particularly a human gene, also is provided. Allelic variants can differ in the identity of a single nucleotide or base pair, for example, by substitution of one nucleotide; in two or more nucleotides or base pairs; or in the number of nucleotides due, for example, to additions or deletions of nucleotides or of trinucleotide repeats; or due to chromosomal rearrangements such as translocations. Specific allelic variants of polymorphic regions are associated with specific diseases and, in some cases, correlate with the prognosis of the disease. A specific allelic variant of a polymorphic region associated with a disease is referred to herein as a "mutant allelic variant" and is considered to be a "genetic lesion."

[0249] Also provided is a process for determining the genetic nature of a phenotype or for identifying a predisposition to that phenotype. For example, it can be determined whether a subject has a predisposition to a specific disease or condition, i.e., whether the subject has, or is at risk of developing, a disease or condition associated with a specific allelic variant of a polymorphic region of a gene. Such a subject can be identified by determining whether the subject carries an allelic variant associated with the specific disease or condition. Furthermore, if the disease is a recessive disease it can be determined whether a subject is a carrier of a recessive allele of a gene associated with the specific disease or condition.

[0250] Numerous diseases or conditions have been genetically linked to a specific gene and, more particularly, to a specific mutation or genetic lesion of a gene. For example, hyperproliferative diseases such as cancers are associated with mutations in specific genes. Such cancers include breast cancer, which has been linked to mutations in BRCA1 or BRCA2. Mutant alleles of BRCA1 are described, for example, in U.S. Patent No. 5,622,829. Other genes such as tumor suppressor genes, which are associated with the development of cancer when mutated, include, but are not limited to, p53 (associated with many forms of cancer); Rb (retinoblastoma); WT1 (Wilm's tumor) and various proto-oncogenes such as c-myc and c-fos (see Thompson and Thompson, "Genetics in Medicine" 5th Ed.; Nora et al., "Medical Genetics" 4th Ed. (Lea and Febiger, eds.)

[0251] A process as disclosed herein also can be used to detect DNA mutations that result in the translation of a truncated polypeptide, as occurs, for example, with BRCA1 and BRCA2. Translation of nucleic acid regions containing such a mutation results in a truncated polypeptide that easily can be differentiated from the corresponding non-truncated polypeptide by mass spectrometry.

[0252] A process as disclosed herein also can be used to genotype a subject, for example, a subject being considered as a recipient or a donor of an organ or a bone marrow graft. For example, the identity of MHC alleles, particularly HLA alleles, in a subject can be determined. The information obtained using such a method is useful because transplantation of a graft to a recipient having different transplantation antigens than the graft can result in rejection of the graft and can result in graft versus host disease following bone marrow transplantation.

[0253] The response of a subject to medicaments can be affected by variations in drug modification systems such as the cytochrome P450 system, and susceptibility to particular infectious diseases can be influenced by genetic status. Thus, the identification of particular allelic variants can be used to predict the potential responsiveness of a subject to specific drug or the susceptibility of a subject to an infectious disease. Genes involved in pharmacogenetics are known (see, e.g., Nora et al., "Medical Genetics" 4th Ed. (Lea and Febiger, eds.).

[0254] Some polymorphic regions may not be related to any disease or condition. For example, many loci in the human genome contain a polymorphic short tandem repeat (STR) region. STR loci contain short, repetitive sequence elements of 3 to 7 base pairs in length. It is estimated that there are 200,000 expected trimeric and tetrameric STRs, which are present as frequently as once every 15 kb in the human genome (see, e.g., International PCT application No. WO 9213969 A1, Edwards et al., Nucl. Acids Res. 19:4791 (1991); Beckmann et al. (1992) Genomics 12:627-631). Nearly half of these STR loci are polymorphic, providing a rich source of genetic markers. Variation in the number of repeat units at a particular locus is responsible for the observed polymorphism reminiscent of variable nucleotide tandem repeat (VNTR) loci (Nakamura et al. (1987) Science 235:1616-1622); and minisatellite loci (Jeffreys et al. (1985) Nature 314: 67-73), which contain longer repeat units, and microsatellite or dinucleotide repeat loci (Luty et al. (1991) Nucleic Acids Res. 19:4308; Litt et al. (1990) Nucleic Acids Res. 18:4301; Litt et al. (1990) Nucleic Acids Res. 18:5921; Luty et al. (1990) Am. J. Hum. Genet. 46:776-783; Tautz (1989) Nucl. Acids Res. 17:6463-6471; Weber et al. (1989) Am. J. Hum. Genet. 44:388-396; Beckmann et al. (1992) Genomics 12:627-631).

[0255] Polymorphic STR loci and other polymorphic regions of genes are extremely useful markers for human iden-

tification, paternity and maternity testing, genetic mapping, immigration and inheritance disputes, zygosity testing in twins, tests for inbreeding in humans, quality control of human cultured cells, identification of human remains, and testing of semen samples, blood stains and other material in forensic medicine. Such loci also are useful markers in commercial animal breeding and pedigree analysis and in commercial plant breeding. Traits of economic importance in plant crops and animals can be identified through linkage analysis using polymorphic DNA markers. Efficient processes for determining the identity of such loci are disclosed herein.

[0256] STR loci can be amplified by PCR using specific primer sequences identified in the regions flanking the tandem repeat to be targeted. Allelic forms of these loci are differentiated by the number of copies of the repeat sequence contained within the amplified region. Examples of STR loci include but are not limited to pentanucleotide repeats in the human CD4 locus (Edwards et al., Nucl. Acids Res. 19:4791 (1991)); tetranucleotide repeats in the human aromatase cytochrome P-450 gene (CYP19; Polymeropoulos et al., Nucl. Acids Res. 19:195 (1991)); tetranucleotide repeats in the human coagulation factor XIII A subunit gene (F13A1; Polymeropoulos et al., Nucl. Acids Res. 19:4306 (1991)); tetranucleotide repeats in the F13B locus (Nishimura et al., Nucl. Acids Res. 20:1167 (1992)); tetranucleotide repeats in the human c-les/fps, proto-oncogene (FES; Polymeropoulos et al., Nucl. Acids Res. 19:4018 (1991)); tetranucleotide repeats in the LFL gene (Zuliani et al., Nucl. Acids Res. 18:4958 (1990)); trinucleotide repeats polymorphism at the human pancreatic phospholipase A-2 gene (PLA2; Polymeropoulos et al., Nucl. Acids Res. 18:7468 (1990)); tetranucleotide repeats polymorphism in the VWF gene (Ploos et al., Nucl. Acids Res. 18:4957 (1990)); and tetranucleotide repeats in the human thyroid peroxidase (hTPO) locus (Anker et al., Hum. Mol. Genet. 1:137 (1992)).

[0257] A target DNA sequence can be part of a foreign genetic sequence such as the genome of an invading microorganism, including, for example, bacteria and their phages, viruses, fungi, protozoa, and the like. The processes provided herein are particularly applicable for distinguishing between different variants or strains of a microorganism in order, for example, to choose an appropriate therapeutic intervention. Examples of disease-causing viruses that infect humans and animals and that can be detected by a disclosed process include but are not limited to *Retroviridae* (e.g., human immunodeficiency viruses such as HIV-1 (also referred to as HTLV-III, LAV or HTLV-III/LAV; Ratner et al., Nature, 313: 227-284 (1985); Wain Hobson et al., Cell, 40:9-17 (1985), HIV-2 (Guyader et al., Nature, 328:662-669 (1987); European Patent Publication No. 0 269 520; Chakrabarti et al., Nature, 328:543-547 (1987); European Patent Application No. 0 655 501), and other isolates such as HIV-LP (International Publication No. WO 94/00562); *Picornaviridae* (e.g., polioviruses, hepatitis A virus, (Gust et al., Intervirology, 20:1-7 (1983)); enteroviruses, human coxsackie viruses, rhinoviruses, echoviruses); *Calciviridae* (e.g. strains that cause gastroenteritis); *Togaviridae* (e.g., equine encephalitis viruses, rubella viruses); *Flaviridae* (e.g., dengue viruses, encephalitis viruses, yellow fever viruses); *Coronaviridae* (e.g., coronaviruses); *Rhabdoviridae* (e.g., vesicular stomatitis viruses, rabies viruses); *Filoviridae* (e.g., ebola viruses); *Paramyxoviridae* (e.g., parainfluenza viruses, mumps virus, measles virus, respiratory syncytial virus); *Orthomyxoviridae* (e.g., influenza viruses); *Bunyaviridae* (e.g., Hantaan viruses, bunya viruses, phleboviruses and Nairo viruses); *Arenaviridae* (hemorrhagic fever viruses); *Reoviridae* (e.g., reoviruses, orbiviruses and rotaviruses); *Birnaviridae*; *Hepadnaviridae* (Hepatitis B virus); *Parvoviridae* (parvoviruses); *Papovaviridae*; *Hepadnaviridae* (Hepatitis B virus); *Parvoviridae* (most adenoviruses); *Papovaviridae* (papilloma viruses, polyoma viruses); *Adenoviridae* (most adenoviruses); *Herpesviridae* (herpes simplex virus type 1 (HSV-1) and HSV-2, varicella zoster virus, cytomegalovirus, herpes viruses); *Poxviridae* (variola viruses, vaccinia viruses, pox viruses); *Iridoviridae* (e.g., African swine fever virus); and unclassified viruses (e.g., the etiological agents of Spongiform encephalopathies, the agent of delta hepatitis (thought to be a defective satellite of hepatitis B virus), the agents of non-A, non-B hepatitis (class 1 = internally transmitted; class 2 = parenterally transmitted, i.e., Hepatitis C); Norwalk and related viruses, and astroviruses.

[0258] Examples of infectious bacteria include but are not limited to *Helicobacter pylori*, *Borelia burgdorferi*, *Legionella pneumophila*, *Mycobacteria* sp. (e.g. *M. tuberculosis*, *M. avium*, *M. intracellulare*, *M. kansaii*, *M. gordonae*), *Staphylococcus aureus*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Listeria monocytogenes*, *Streptococcus pyogenes* (Group A Streptococcus), *Streptococcus agalactiae* (Group B Streptococcus), *Streptococcus* sp. (viridans group), *Streptococcus faecalis*, *Streptococcus bovis*, *Streptococcus* sp. (anaerobic species), *Streptococcus pneumoniae*, pathogenic *Campylobacter* sp., *Enterococcus* sp., *Haemophilus influenzae*, *Bacillus anthracis*, *Corynebacterium diphtheriae*, *Corynebacterium* sp., *Erysipelothrix rhusiopathiae*, *Clostridium perfringens*, *Clostridium tetani*, *Enterobacter aerogenes*, *Klebsiella pneumoniae*, *Pasturella multocida*, *Bacteroides* sp., *Fusobacterium nucleatum*, *Streptobacillus moniliformis*, *Treponema pallidum*, *Treponema pertenue*, *Leptospira*, and *Actinomyces israeli*.

[0259] Examples of infectious fungi include but are not limited to *Cryptococcus neoformans*, *Histoplasma capsulatum*, *Coccidioides immitis*, *Blastomyces dermatitidis*, *Chlamydia trachomatis*, *Candida albicans*. Other infectious organisms include protists such as *Plasmodium falciparum* and *Toxoplasma gondii*.

[0260] The processes provided herein are further illustrated by the following examples, which should not be construed as limiting in any way. The practice of the processes will employ, unless otherwise indicated, conventional techniques of cell biology, cell culture, molecular biology, transgenic biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature. See, for example, DNA Cloning, Volumes I and II (D.N. Glover ed., 1985); Oligonucleotide Synthesis (M.J. Gait ed., 1984); Mullis et al. U.S. Patent No:

4,683,194; Nucleic Acid Hybridization (B.D. Hames & S.J. Higgins eds. 1984); Transcription and Translation (B.D. Hames & S.J. Higgins eds. 1984); Culture of Animal Cells (R.I. Freshney, Alan R. Liss, Inc., 1987); Immobilized Cells and Enzymes (IRL Press, 1986); B. Perbal, A Practical Guide to Molecular Cloning (1984); the treatise, Methods In Enzymology (Academic Press, Inc., N.Y.); Gene Transfer Vectors For Mammalian Cells (J.H. Miller and M.P. Calos eds., 1987, Cold Spring Harbor Laboratory); Methods In Enzymology, Vols. 154 and 155 (Wu et al. eds.), Immunochemical Methods In Cell And Molecular Biology (Mayer and Walker, eds., Academic Press, London, 1987); Handbook Of Experimental Immunology, Volumes I-IV (D. M. Weir and C.C. Blackwell, eds., 1986); Manipulating the Mouse Embryo (Cold Spring Harbor Laboratory press, Cold Spring Harbor, N.Y., 1986).

[0261] The following examples are included for illustrative purposes only and are not intended to limit the scope of the invention.

EXAMPLE 1

[0262] This example which is not within the scope of the invention demonstrates that genomic DNA obtained from patients with spinal cerebellar ataxia 1 (SCA-1) can be used to identify target polypeptides encoded by trinucleotide repeats associated with SCA-1.

Genomic DNA Amplification

[0263] Human genomic DNA was extracted using the QIAMP Blood Kit (Qiagen), following the manufacturer's protocol. A region of the extracted DNA containing the (CAG) repeat associated with SCA-1 was amplified by PCR using primers modified to contain a transcription promoter sequence and a region coding for a His-6 tag peptide. The forward primer had the following nucleotide sequence, in which the T7 promoter sequence is italicized and the bases on the 5'-side of the promoter are random:

**5'-d(GAC TTT ACT TGT ACG TGC ATA ATA CGA CTC ACT ATA GGG
AGA CTG ACC ATG GGC AGT CTG AGC CA) (SEQ ID NO: 6).**

[0264] The reverse primer had the following nucleotide sequence, in which the nucleotide sequence encoding the His-6 tag peptide is represented in bold and the first six 5'-bases are random:

**5'-d(TGA TTC TCA ATG ATG ATG ATG ATG AAC TTG AAA TGT
GGA CGT AC) (SEQ ID NO: 7).**

[0265] Total reaction volume was 50 µl with 20 pmol primers per reaction. Taq polymerase including 10X buffer was obtained from Boehringer Mannheim and dNTPs were obtained from Pharmacia. Cycling conditions included 5 min at 94°C, followed by 35 cycles of 30 sec at 94°C, 45 sec at 53°C, 30 sec at 72°C, with a final extension time of 2 min at 72°C. PCR products were purified using the Qiagen QIAQUICK kit and elution of the purified products was performed using 50 µL 10 mM Tris-HCl buffer (pH 8).

Coupled *In Vitro* Transcription and Translation

[0266] Coupled transcription and translation was performed using the TNT reaction buffer (Promega). Reaction components, in a total volume of 50 µl. were thawed and mixed according to the manufacturer's protocol, using 1 µl of T7 RNA polymerase and 1 pmol of amplified DNA, except that unlabeled methionine was used in place of ³⁵S-methionine. The reaction mixture was incubated at 30°C for 90 min.

Target Polypeptide Purification

[0267] The translated His-6 tagged polypeptide was purified from the wheat germ extract mixture using the Qiagen QIAEXPRESS Ni-NTA protein purification system according to the manufacturer's protocol. Briefly, the extract mixture was washed by centrifugation through a spin column containing a nickel-nitriloacetic acid resin, which affinity captures the His-6 peptide tag on the polypeptide. The polypeptide was eluted from the column with 100 mM imidazole.

Mass Spectrometry

[0268] The translated polypeptide was mixed with matrix either directly from the elution solution or first was lyophilized and resuspended in 5 μ l H₂O. This solution was mixed 1:1 (v:v) with matrix solution (concentrated sinnapinic acid in 50/50 v:v ethanol/H₂O), and 0.5 μ l of the mixture was added to a sample probe for analysis in a linear time-of-flight mass spectrometer operated in delayed ion extraction mode with a source potential of 25 kV. Internal calibration was achieved for all spectra using three intense matrix ion signals.

RESULTS

[0269] Genomic DNA was obtained from 4 patients having SCA-1, as described above. Three of the patients had 10, 15, or 16 CAG repeats and the fourth patient had an unknown number of trinucleotide repeats.

[0270] A region containing the trinucleotide repeats was PCR amplified using primers (SEQ ID NOS: 6 and 7) that hybridized to sequences located on either side of the repeats. The nucleotide sequence (SEQ ID NO: 8) of a PCR product amplified from a region containing 10 CAG repeats is shown in Figure 1A and the amino acid sequence (SEQ ID NO: 9) of a polypeptide encoded by the amplified nucleic acid is shown in Figure 1B (SEQ ID NO: 9).

[0271] The amplified DNA from each patient was subjected to *in vitro* transcription and translation, and the target polypeptides were isolated on a nickel chromatography column. Mass spectrometric analysis of the peptides encoded by target polypeptides encoded by the 10, 15, and 16 CAG repeats indicated that these peptides had a molecular mass of 8238.8, 8865.4, and 8993.6 Daltons, respectively. The polypeptide encoded by the nucleic acid from the fourth patient, having an unknown number of trinucleotide repeats, had a molecular weight of 8224.8 Da. While this value does not correspond exactly with a unit number of repeats (10 is the closest), it is consistent with detection of a point mutation; i.e., the -14 Dalton shift for this polypeptide corresponds to an Ala-> Gly mutation due to a C-> G mutation in one of the repeats. This result demonstrates that the disclosed process allows the identification of a target polypeptide encoded by a genetic lesion associated with a disease. In addition, the results demonstrate that such a process allows the detection of a single base difference between two nucleic acids.

[0272] Detection of such subtle differences in the protein lengths are not reproducibly obtained with electrophoretic methods even with use of multiple internal standards. Even low performance MS instrumentation is capable of far better than 0.1 % mass accuracy in this mass range using internal calibration; higher performance instrumentation such as Fourier transform MS is capable of ppm mass accuracy with internal or external calibration. It should be noted that the mass difference between the 15 and 16 repeat unit polypeptides is 1.4% and the 14 Dalton mass shift due to the point mutation between the 10 repeat patients is 0.17%. Clearly, each of these situations can be routinely analyzed successfully.

EXAMPLE 2

1-(2-Nitro-5-(3-O-4,4'-dimethoxytritylpropoxy)phenyl)-1-O-((2-cyanoethoxy)-diisopropylaminophosphino) ethane

A. 2-Nitro-5-(3-hydroxypropoxy)benzaldehyde

[0273] 3-Bromo-1-propanol (3.34 g, 24 mmol) was refluxed in 80 ml of anhydrous acetonitrile with 5-hydroxy-2-nitrobenzaldehyde (3.34 g, 20 mmol), K₂CO₃ (3.5 g), and KI (100 mg) overnight (15 hr). The reaction mixture was cooled to room temperature and 150 ml of methylene chloride was added. The mixture was filtered and the solid residue was washed with methylene chloride. The combined organic solution was evaporated to dryness and redissolved in 100 ml methylene chloride. The resulted solution was washed with saturated NaCl solution and dried over sodium sulfate. 4.31 g (96%) of desired product was obtained after removal of the solvent *in vacuo*.

R_f = 0.33 (dichloromethane/methanol, 95/5).

UV (methanol) maximum: 313, 240 (shoulder), 215 nm; minimum: 266 nm.

¹H NMR (DMSO-d₆) δ 10.28 (s, 1H), 8.17 (d, 1H), 7.35 (d, 1H), 7.22 (s, 1H), 4.22 (t, 2H), 3.54 (t, 2H), 1.90 (m, 2H).

¹³C NMR (DMSO-d₆) δ 189.9, 153.0, 141.6, 134.3, 127.3, 118.4, 114.0, 66.2, 56.9, 31.7.

B. 2-Nitro 5-(3-O-t-butylidimethylsilylpropoxy)benzaldehyde

[0274] 2-Nitro-5-(3-hydroxypropoxy)benzaldehyde (1 g, 4.44 mmol) was dissolved in 50 ml anhydrous acetonitrile. To this solution was added 1 ml of triethylamine, 200 mg of imidazole, and 0.8 g (5.3 mmol) of TBDMSCl. The mixture was stirred at room temperature for 4 hr. Methanol (1 ml) was added to stop the reaction. The solvent was removed *in vacuo* and the solid residue was redissolved in 100 ml methylene chloride. The resulting solution was washed with saturated

sodium bicarbonate solution and then water.' The organic phase was dried over sodium sulfate and the solvent was removed *in vacuo*. The crude mixture was subjected to a quick silica gel column with methylene chloride to yield 1.44 g (96%) of 2-nitro-5-(3-O-t-butyl dimethylsilylpropoxy)benzaldehyde.

$R_f = 0.67$ (hexane/ethyl acetate, 5/1).

5 UV (methanol), maximum: 317, 243, 215 nm; minimum: 235, 267 nm.

^1H NMR (DMSO- d_6) δ 10.28 (s, 1H), 8.14 (d, 1H), 7.32 (d, 1H), 7.20 (s, 1H), 4.20 (t, 2H), 3.75 (t, 2H), 1.90 (m, 2H), 0.85 (s, 9H), 0.02 (s, 6H). ^{13}C NMR (DMSO- d_6) δ 189.6, 162.7, 141.5, 134.0, 127.1, 118.2, 113.8, 65.4, 58.5, 31.2, 25.5, -3.1, -5.7.

10 C. 1-(2-Nitro-5-(3-O-t-butyl dimethylsilylpropoxy)phenyl)ethanol

[0275] High vacuum dried 2-nitro-5-(3-O-t-butyl dimethylsilylpropoxy) benzaldehyde (1.02 g, 3 mmol) was dissolved 50 ml of anhydrous methylene chloride. 2 M trimethylaluminium in toluene (3 ml) was added dropwise within 10 min and kept the reaction mixture at room temperature. It was stirred further for 10 min and the mixture was poured into 10 ml ice cooled water. The emulsion was separated from water phase and dried over 100 g of sodium sulfate to remove the remaining water. The solvent was removed *in vacuo* and the mixture was applied to a silica gel column with gradient methanol in methylene chloride. 0.94 g (86%) of desired product was isolated.

$R_f = 0.375$ (hexane/ethyl acetate, 5/1)

UV (methanol), maximum: 306, 233, 206 nm; minimum: 255, 220 nm.

20 ^1H NMR (DMSO- d_6) δ 8.00 (d, 1H), 7.36 (s, 1H), 7.00 (d, 1H), 5.49 (b, OH), 5.31 (q, 1H), 4.19 (m, 2H), 3.77 (t, 2H), 1.95 (m, 2H), 1.37 (d, 3H), 0.86 (s, 9H), 0.04 (s, 6H).

^{13}C NMR (DMSO- d_6) δ 162.6, 146.2, 139.6, 126.9, 112.9, 112.5, 64.8, 63.9, 58.7, 31.5, 25.6, 24.9, -3.4, -5.8

25 D. 1-(2-Nitro-5-(3-hydroxypropoxy)phenyl)ethanol

[0276] 1-(2-Nitro-5-(3-O-t-butyl dimethylsilylpropoxy)phenyl)ethanol (0.89 g, 2.5 mmol) was dissolved in 30 ml of THF and 0.5 mmol of $n\text{Bu}_4\text{NF}$ was added under stirring. The mixture was stirred at room temperature for 5 hr and the solvent was removed *in vacuo*. The remaining residue was applied to a silica gel column with gradient methanol in methylene chloride. 1-(2-nitro-5-(3-hydroxypropoxy)phenyl)ethanol (0.6 g (99%)) was obtained.

30 $R_f = 0.17$ (dichloromethane/methanol, 95/5).

UV (methanol), maximum: 304, 232, 210 nm; minimum: 255, 219 nm.

^1H NMR (DMSO- d_6) δ 8.00 (d, 1H), 7.33 (s, 1H), 7.00 (d, 1H), 5.50 (d, OH), 5.28 (t, OH), 4.59 (t, 1H), 4.17 (t, 2H), 3.57 (m, 2H), 1.89 (m, 2H), 1.36 (d, 2H).

^{13}C NMR (DMOS- d_6) δ 162.8, 146.3, 139.7, 127.1, 113.1, 112.6, 65.5, 64.0, 57.0, 31.8, 25.0

35 E. 1-(2-Nitro-5-(3-O-4,4'-dimethoxytritylpropoxy)phenyl)ethanol

[0277] 1-(2-Nitro-5-(3-hydroxypropoxy)phenyl)ethanol (0.482 g, 2 mmol) was co-evaporated with anhydrous pyridine twice and dissolved in 20 ml anhydrous pyridine. The solution was cooled in ice water bath and 750 mg (2.2 mmol) of DMTCI was added. The reaction mixture was stirred at room temperature overnight and 0.5 ml methanol was added to stop the reaction. The solvent was removed *in vacuo* and the residue was co-evaporated with toluene twice to remove trace of pyridine. The final residue was applied to a silica gel column with gradient methanol in methylene chloride containing drops of triethylamine to yield 0.96 g (89%) of the desired product 1-(2-nitro-5-(3-O-4,4'-dimethoxytritylpropoxy)phenyl)ethanol.

45 $R_f = 0.50$ (dichloromethane/methanol, 99/1)

UV (methanol), maximum: 350 (shoulder), 305, 283, 276 (shoulder), 233, 208 nm; minimum: 290, 258, 220 nm.

^1H NMR (DMSO- d_6) δ 8.00 (d, 1H), 6.82-7.42 (ArH), 5.52 (d, OH), 5.32 (m, 1H), 4.23 (t, 2H), 3.71 (s, 6H), 3.17 (t, 2H), 2.00 (m, 2H), 1.37 (d, 3H).

50 ^{13}C NMR (DMOS- d_6) δ 162.5, 157.9, 157.7, 146.1, 144.9, 140.1, 139.7, 135.7, 129.5, 128.8, 127.6, 127.5, 127.3, 126.9, 126.4, 113.0, 112.8, 112.6, 85.2, 65.3, 63.9, 59.0, 54.8, 28.9, 24.9.

F. 1-(2-Nitro-5-(3-O-4,4'-dimethoxytritylpropoxy)phenyl)-1-O-(12-cyanoethoxy)-diisopropylaminophosphino ethane

55 [0278] 1-(2-Nitro-5-(3-O-4,4'-dimethoxytritylpropoxy)phenyl)ethanol (400 mg, 0.74 mmol) was dried under high vacuum and was dissolved in 20 ml of anhydrous methylene chloride. To this solution, it was added 0.5 ml N,N-diisopropylethylamine and 0.3 ml (1.34 mmol) of 2-cyanoethyl-N,N-diisopropylchlorophosphoramidite. The reaction mixture was stirred at room temperature for 30 min and 0.5 ml of methanol was added to stop the reaction. The mixture was washed

with saturated sodium bicarbonate solution and was dried over sodium sulfate. The solvent was removed *in vacuo* and a quick silica gel column with 1 % methanol in methylene chloride containing drops of triethylamine yield 510 mg (93%) the desired phosphoramidite. $R_f=0.87$ (dichloromethane/methanol, 99/11).

5 EXAMPLE 3

1-(4-(3-O-4,4'-Dimethoxytritylpropoxy)-3-methoxy-6-nitrophenyl)-1-O-((2-cyanoethoxy)-diisopropylamino-phosphino)ethane

10 A. 4-(3-Hydroxypropoxy)-3-methoxyacetophenone

[0279] 3-Bromo-1-propanol (53 ml, 33 mmol) was refluxed in 100 ml of anhydrous acetonitrile with 4-hydroxy-3-methoxyacetophenone (5 g, 30 mmol), K_2CO_3 (5 g), and KI (300 mg) overnight (15 h). Methylene chloride (150 ml) was added to the reaction mixture after cooling to room temperature. The mixture was filtered and the solid residue was washed with methylene chloride. The combined organic solution was evaporated to dryness and redissolved in 100 ml methylene chloride. The resulted solution was washed with saturated NaCl solution and dried over sodium sulfate. 6.5 g (96.4%) of desired product was obtained after removal of the solvent *in vacuo*.

$R_f = 0.41$ (dichloromethane/methanol, 95/5)

UV (methanol), maximum: 304, 273, 227, 210 nm; minimum: 291, 244, 214 nm.

1H NMR (DMSO- d_6) δ 7.64 (d, 1 H), 7.46 (s, 1 H), 7.04 (d, 1H), 4.58 (b, OH), 4.12 (t, 2H), 3.80 (s; 3H), 3.56 (t, 2H), 2.54 (s, 3H), 1.88 (m, 2H).

^{13}C NMR (DMSO- d_6) δ 196.3, 152.5, 148.6, 129.7, 123.1, 111.5, 110.3, 65.4, 57.2, 55.5, 31.9, 26.3.

25 B. 4-(3-Acetoxypropoxy)-3-methoxyacetophenone

[0280] 4-(3-Hydroxypropoxy)-3-methoxyacetophenone (3.5 g, 15.6 mmol) was dried and dissolved in 80 ml anhydrous acetonitrile. To this mixture, 6 ml of triethylamine and 6 ml of acetic anhydride were added. After 4 h, 6 ml methanol was added and the solvent was removed *in vacuo*. The residue was dissolved in 100 ml dichloromethane and the solution was washed with dilute sodium bicarbonate solution, then water. The organic phase was dried over sodium sulfate and the solvent was removed. The solid residue was applied to a silica gel column with methylene chloride to yield 4.1 g of 4-(3-acetoxypropoxy)-3-methoxyacetophenone (98.6%).

$R_f = 0.22$ (dichloromethane/methanol, 99/1).

UV (methanol), maximum: 303, 273, 227, 210 nm; minimum: 290, 243, 214 nm.

1H NMR (DMSO- d_6) δ 7.62 (d, 1H), 7.45 (s, 1H), 7.08 (d, 1H), 4.12 (m, 4H), 3.82 (s, 3H), 2.54 (s, 3H), 2.04 (m, 2H), 2.00 (s, 3H).

^{13}C NMR (DMSO- d_6) δ 196.3, 170.4, 152.2, 148.6, 130.0, 123.0, 111.8, 110.4, 65.2, 60.8, 55.5, 27.9, 26.3, 20.7.

C. 4-(3-Acetoxypropoxy)-3-methoxy-6-nitroacetophenone

[0281] 4-(3-Acetoxypropoxy)-3-methoxyacetophenone (3.99 g, 15 mmol) was added portionwise to 15 ml of 70% HNO_3 in water bath; the reaction temperature was maintained at the room temperature. The reaction mixture was stirred at room temperature for 30 min and 30 g of crushed ice was added. This mixture was extracted with 100 ml of dichloromethane and the organic phase was washed with saturated sodium bicarbonate solution. The solution was dried over sodium sulfate and the solvent was removed *in vacuo*. The crude mixture was applied to a silica gel column with gradient methanol in methylene chloride to yield 3.8 g (81.5%) of desired product 4-(3-acetoxypropoxy)-3-methoxy-6-nitroacetophenone and 0.38 g (8%) of ipso-substituted product 5-(3-acetoxypropoxy)-4-methoxy-1,2-dinitrobenzene. Side ipso-substituted product 5-(3-acetoxypropoxy)-4-methoxy-1,2-dinitrobenzene:

$R_f = 0.47$ (dichloromethane/methanol, 99/1).

UV (methanol), maximum: 334, 330, 270, 240, 212 nm; minimum: 310, 282, 263, 223 nm.

1H NMR ($CDCl_3$) δ 7.36 (s, 1H), 7.34 (s, 1H), 4.28 (t, 2H), 4.18 (t, 2H), 4.02 (s, 3H), 2.20 (m, 2H), 2.08 (s, 3H).

^{13}C NMR ($CDCl_3$) δ 170.9, 152.2, 151.1, 117.6, 111.2, 107.9, 107.1, 66.7, 60.6, 56.9, 28.2, 20.9.

Desired product 4-(3-acetoxypropoxy)-3-methoxy-6-nitroacetophenone:

$R_f=0.29$ (dichloromethane/methanol, 99/1).

UV (methanol), maximum: 344, 300, 246, 213 nm; minimum: 320, 270, 227 nm.

1H NMR ($CDCl_3$) δ 7.62 (s, 1H), 6.74 (s, 1H), 4.28 (t, 2H), 4.20 (t, 2H), 3.96 (s, 3H), 2.48 (s, 3H), 2.20 (m, 2H), 2.08 (s, 3H).

^{13}C NMR ($CDCl_3$) δ 200.0, 171.0, 154.3, 148.8, 138.3, 133.0, 108.8, 108.0, 66.1, 60.8, 56.6, 30.4, 28.2, 20.9.

D. 1-(4-(3-Hydroxypropoxy)-3-methoxy-6-nitrophenyl)ethanol

[0282] 4-(3-Acetoxypropoxy)-3-methoxy-6-nitroacetophenone (3.73 g, 12 mmol) was added 150 ml ethanol and 6.5 g of K_2CO_3 . The mixture was stirred at room temperature for 4 hr and TLC with 5% methanol in dichloromethane indicated the completion of the reaction. To this same reaction mixture was added 3.5 g of $NaBH_4$ and the mixture was stirred at room temperature for 2 hr. Acetone (10 ml) was added to react with the remaining $NaBH_4$. The solvent was removed *in vacuo* and the residue was uptaken into 50 g of silica gel. The silica gel mixture was applied on the top of a silica gel column with 5% methanol in methylene chloride to yield 3.15 g (97%) of desired product 1-(4-(3-hydroxypropoxy)-3-methoxy-6-nitrophenyl)ethanol. Intermediate product 4-(3-hydroxypropoxy)-3-methoxy-6-nitroacetophenone after de-

protection:

$R_f=0.60$ (dichloromethane/methanol, 95/5).

Final product 1-(4-(3-hydroxypropoxy)-3-methoxy-6-nitrophenyl)ethanol:

$R_f=0.50$ (dichloromethane/methanol, 95/5).

UV (methanol), maximum: 344, 300, 243, 219 nm; minimum: 317, 264, 233 nm.

1H NMR ($DMSO-d_6$) δ 7.54 (s, 1H), 7.36 (s, 1H), 5.47 (d, OH), 5.27 (m, 1H), 4.55 (t, OH), 4.05 (t, 2H), 3.90 (s, 3H), 3.55 (q, 2H), 1.88 (m, 2H), 1.37 (d, 3H).

^{13}C NMR ($DMSO-d_6$) δ 153.4, 146.4, 138.8, 137.9, 109.0, 108.1, 68.5, 65.9, 57.2, 56.0, 31.9, 29.6.

E. 1-(4-(3-O-4,4'-Dimethoxytritylpropoxy)-3-methoxy-6-nitrophenyl)ethanol

[0283] 1-(4-(3-Hydroxypropoxy)-3-methoxy-6-nitrophenyl)ethanol (0.325 g, 1.2 mmol) was co-evaporated with anhydrous pyridine twice and dissolved in 15 ml anhydrous pyridine. The solution was cooled in ice-water bath and 450 mg (1.33 mmol) of DMTCl was added. The reaction mixture was stirred at room temperature overnight and 0.5 ml methanol was added to stop the reaction. The solvent was removed *in vacuo* and the residue was co-evaporated with toluene twice to remove trace of pyridine. The final residue was applied to a silica gel column with gradient methanol in methylene chloride containing drops of triethylamine to yield 605 mg (88%) of desired product 1-(4-(3-O-4,4'-dimethoxytritylpropoxy)-3-methoxy-6-nitrophenyl)ethanol.

$R_f=0.50$ (dichloromethane/methanol, 95/5).

UV (methanol), maximum: 354, 302, 282, 274, 233, 209 nm; minimum: 322, 292, 263, 222 nm.

1H NMR ($DMSO-d_6$) δ 7.54 (s, 1 H), 6.8-7.4 (ArH), 5.48 (d, OH), 5.27 (m, 1 H), 4.16 (t, 2H), 3.85 (s, 3H), 3.72 (s, 6H), 3.15 (t, 2H), 1.98 (t, 2H), 1.37 (d, 3H).

^{13}C NMR ($DMSO-d_6$) δ 157.8, 153.3, 146.1, 144.9, 138.7, 137.8, 135.7, 129.4, 128.7, 127.5, 127.4, 126.3, 112.9, 112.6, 108.9, 108.2, 85.1, 65.7, 63.7, 59.2, 55.8, 54.8, 29.0, 25.0.

F. 1-(4-(3-O-4,4'-Dimethoxytritylpropoxy)-3-methoxy-6-nitrophenyl)-1-O-((2-cyanoethoxy)-diisopropylamino-phosphino)ethane

[0284] 1-(4-(3-O-4,4'-Dimethoxytritylpropoxy)-3-methoxy-6-nitrophenyl)ethanol (200 mg, 3.5 mmol) was dried under high vacuum and was dissolved in 15 ml of anhydrous methylene chloride. To this solution, it was added 0.5 ml N,N-diisopropylethylamine and 0.2 ml (0.89 mmol) of 2-cyanoethyl-N,N-diisopropylchlorophosphoramidite. The reaction mixture was stirred at room temperature for 30 min and 0.5 ml of methanol was added to stop the reaction. The mixture was washed with saturated sodium bicarbonate solution and was dried over sodium sulfate. The solvent was removed *in vacuo* and a quick silica gel column with 1 % methanol in methylene chloride containing drops of triethylamine yield 247 mg (91.3%) the desired phosphoramidite 1-(4-(3-O-4,4'-dimethoxytritylpropoxy)-3-methoxy-6-nitrophenyl)-1-O-((2-cyanoethoxy)-diisopropylaminophosphino)ethan. $R_f = 0.87$ (dichloromethane/methanol, 99/1)

SEQUENCE LISTING

[0285]

<110> Little, Daniel
Higgins, G. Scott
Koster, Hubert
Lough, David
SEQUENOM, INC.

<120> Mass Spectrometric Detection of Polypeptides

<130> 2016B
 <140> Unassigned
 <141> 1998-09-02
 5
 <150> 08/922,201
 <151> 1997-09-02
 <160>
 10
 <170> PatentIn Ver. 2.0
 <210> 1
 <211> 24
 15
 <212> DNA
 <213> Bacteriophage SP6
 <220>
 <221> promoter
 20
 <222> (1)..(24)
 <223> SP6 promoter sequence (single-stranded)
 <400> 1
 25
 catacgattt aggtgacact atag 24
 <210> 2
 <211> 18
 <212> DNA
 <213> Bacteriophage SP6
 30
 <220>
 <221> promoter
 <222> (1) .. (18)
 <223> SP6 promoter sequence (single-stranded)
 35
 <400> 2
 atttagtga cactatag 18
 <210> 3
 40
 <211> 20
 <212> DNA
 <213> Bacteriophage T3
 <220>
 45
 <221> promoter
 <222> (1)..(2-0)
 <223> T3 promoter sequence (single-stranded)
 <400> 3
 50
 attaacccctc actaaaggga 20
 <210> 4
 <211> 20
 <212> DNA
 55
 <213> Bacteriophage T7
 <220>
 <221> promoter

<222> (1)..(20)
<223> T7 promoter sequence (single-stranded)

5 <400> 4
taatacgact cactataggg 20

<210> 5
<211> 8
<212> DNA
10 <213> Prokaryote

<220>
<221> misc_feature
<222> (1) .. (8)
15 <223> Primer sequence containing the Shine-Dalgarno (prokaryotic ribosome binding) sequence

<400> 5
taaggagg 8
20 <210> 6
<211> 65
<212> DNA
<213> Artificial Sequence

25 <220>
<223> Description of Artificial Sequence: Primer containing T7 promoter sequence

<220>
<221> promoter
30 <222> (19)..(42)
<223> T7 promoter sequence located within primer as indicated

<400> 6
35 gactttactt gtacgtgcat aatacgactc actatagggga gactgaccat gggcagtcctg 60
agcca 65

40 <210> 7
<211> 47
<212> DNA
<213> Artificial Sequence

45 <220>
<223> Description of Artificial Sequence: Primer encoding His-6 "tag" peptide

<220>
<221> repeat_region
50 <222> (10)..(27)
<223> Sequence encoding His-6 "tag" feature located within primer as indicated

<400> 7
55 tgattctcaa tgatgatgat gatgatgaac ttgaaatgtg gacgtac 47
<210> 8
<211> 270
<212> DNA

<213> Homo sapiens

<220>

<221> repeat_region

5 <222> (8)..(162)

<223> "CAG" repeat region associated with spinal cerebellar ataxia 1 (SCA-1)

<220>

<221> repeat_region

10 <222> (244)..(261)

<223> His-6 "tag" region

<400> 8

15

gactttactt gtacgtgcat aatacgactc actataggga gactgaac 48

atg ggc agt ctg agc cag acg ccg gga cac aag gct gag cag cag cag 96
Met Gly Ser Leu Ser Gln Thr Pro Gly His Lys Ala Glu Gln Gln Gln

20

cag cag cag cag cag cag cag cag cag cat cag cat cag cag cag cag 144
Gln Gln Gln Gln Gln Gln Gln Gln Gln His Gln His Gln Gln Gln Gln

cag cag cag cag cag cag cac ctc acg agg gct ccg ggc ctc atc acc 192
Gln Gln Gln Gln Gln Gln His Leu Ser Arg Ala Pro Gly Leu Ile Thr

25

ccg ggt ccc ccc cac cag ccc agc aga acc agt acg tcc aca ttt caa 240
Pro Gly Pro Pro Gly Gln Pro Ser Arg Thr Ser Thr Ser Thr Gly Gln

gtt cat cat cat cat cat cat cat tgagaatca 270
Val His His His His His His

30

<210> 9

<211> 71

<212> PRT

35

<213> Homo sapiens

<220>

<221> REPEAT

<222> (14)..(38)

40

<223> "Gln" repeat region associated with spinal cerebellar ataxia 1 (SCA-1)

<220>

<221> REPEAT

<222> (66)..(71)

45

<223> His-6 "tag"

<400> 9

50

55

Met Gly Ser Leu Ser Gln Thr Pro Gly His Lys Ala Glu Gln Gln Gln
1 5 10 15
5
Gln Gln Gln Gln Gln Gln Gln Gln Gln His Gln His Gln Gln Gln Gln
20 25 30
10
Gln Gln Gln Gln Gln Gln His Leu Ser Arg Ala Pro Gly Leu Ile Thr
35 40 45
15
Pro Gly Pro Pro Gly Gln Pro Ser Arg Thr Ser Thr Ser Thr Gly Gln
50 55 60
20
Val His His His His His His
65 70
25

Claims

1. A process for determining the amino acid sequence of each polypeptide in a plurality of polypeptides using mass spectroscopy, comprising the steps of:
 - a) differentially mass modifying each polypeptide in the plurality to produce differentially mass modified polypeptides;
 - b) contacting the differentially mass modified polypeptides with an agent that cleaves an amino acid from a terminus of the polypeptides to produce a cleaved amino acid and a deletion fragment;
 - c) subjecting the cleaved amino acid or the deletion fragment to mass spectrometry; and
 - d) repeating step b) and step c), as necessary, thereby determining the amino acid sequence of each polypeptide in the plurality.
2. The process of claim 1, wherein each polypeptide in the plurality is immobilized to the solid support.
3. The process of claim 2, wherein each polypeptide in the plurality is immobilized to the solid support through a cleavable linker.
4. The process of claim 3, wherein the cleavable linker is selected from the group consisting of an acid cleavable linker and photocleavable linker.
5. The process of claim 1, further comprising conditioning each polypeptide prior to step b), or conditioning the cleaved amino acid or the deletion fragment prior to mass spectrometry.
6. The process of claim 1, wherein the conditioning comprises contacting the target polypeptide with a cation exchange material.
7. The process of claim 1, wherein the agent is a chemical agent.
8. The process of claim 1, wherein the agent is an enzyme.

9. The process of claim 8, wherein the enzyme is an aminopeptidase or a carboxypeptidases.
10. The process of claim 1, wherein each polypeptide in the plurality is immobilized on a solid support.
- 5 11. The process of claim 10, wherein each polypeptide is immobilized in an array.

Patentansprüche

- 10 1. Verfahren zur Bestimmung der Aminosäuresequenz eines jeden Polypeptids in einer Vielzahl von Polypeptiden mittels Massenspektroskopie, umfassend die Schritte:
 - a) Differenzielles Modifizieren der Masse jedes Polypeptids in der Vielzahl um Polypeptide mit differenziell modifizierten Massen zu erzeugen;
 - 15 b) Inkontaktbringen der Polypeptide mit differenziell modifizierten Massen mit einem Mittel, das eine Aminosäure von einem Terminus der Polypeptide abspaltet um eine abgespaltene Aminosäure und ein Deletionsfragment zu erzeugen;
 - c) Unterwerfen der abgespaltenen Aminosäure oder des Deletionsfragmentes der Massenspektrometrie; und
 - 20 d) Wiederholen von Schritt b) und von Schritt c) nach Bedarf, und hierdurch Bestimmen der Aminosäuresequenz eines jeden Polypeptids in der Vielzahl.
2. Verfahren nach Anspruch 1, wobei jedes Polypeptid in der Vielzahl an dem festen . Träger immobilisiert ist.
3. Verfahren nach Anspruch 2, wobei jedes Polypeptid in der Vielzahl durch einen spaltbaren Linker an dem festen
25 Träger immobilisiert ist.
4. Verfahren nach Anspruch 3, wobei der spaltbare Linker aus der Gruppe, bestehend aus einem durch Säure spaltbaren Linker und einem durch Licht spaltbaren Linker, ausgewählt wird.
- 30 5. Verfahren nach Anspruch 1, weiterhin umfassend das Konditionieren jedes Polypeptids vor Schritt b) oder das Konditionieren der abgespaltenen Aminosäure oder des Deletionsfragmentes vor der Massenspektrometrie.
6. Verfahren nach Anspruch 1, wobei das Konditionieren das Inkontaktbringen des Ziel-Polypeptids mit einem Kationenaustauschermaterial umfasst.
- 35 7. Verfahren nach Anspruch 1, wobei das Mittel ein chemisches Mittel ist.
8. Verfahren nach Anspruch 1, wobei das Mittel ein Enzym ist.
- 40 9. Verfahren nach Anspruch 8, wobei das Enzym eine Aminopeptidase oder eine Carboxypeptidase ist.
10. Verfahren nach Anspruch 1, wobei jedes Polypeptid in der Vielzahl auf einem festen Träger immobilisiert ist.
11. Verfahren nach Anspruch 10, wobei jedes Polypeptid in einem Array immobilisiert ist.

Revendications

- 50 1. Procédé pour déterminer la séquence d'acides aminés de chaque polypeptide dans une pluralité de polypeptides en utilisant la spectroscopie de masse, comprenant les étapes consistant à :
 - a) modifier de façon différentielle la masse de chaque polypeptide de la pluralité pour produire des polypeptides à masse modifiée de façon différentielle ;
 - 55 b) mettre en contact les polypeptides à masse modifiée de façon différentielle avec un agent qui clive un acide aminé d'une extrémité terminale des polypeptides pour produire un acide aminé clivé et un fragment de délétion clivé ; et
 - c) soumettre l'acide aminé clivé ou le fragment de délétion à la spectrométrie de masse ; et
 - d) répéter l'étape b) et l'étape c), autant que nécessaire, pour déterminer ainsi la séquence d'acides aminés

de chaque polypeptide de la pluralité.

2. Procédé selon la revendication 1, dans lequel chaque polypeptide de la pluralité est immobilisé sur le support solide.

5 3. Procédé selon la revendication 2, dans lequel chaque polypeptide est immobilisé sur le support solide par un agent de liaison clivable.

4. Procédé selon la revendication 3, dans lequel l'agent de liaison clivable est choisi dans le groupe formé par un agent de liaison clivable par acide et un agent de liaison photoclivable.

10

5. Procédé selon la revendication 1, comprenant en outre l'étape consistant à conditionner chaque polypeptide avant l'étape b), ou à conditionner l'acide aminé clivé ou le fragment de délétion avant la spectrométrie de masse.

15

6. Procédé selon la revendication 1, dans lequel le conditionnement comprend la mise en contact du polypeptide cible avec un matériau à échange de cation.

7. Procédé selon la revendication 1, dans lequel l'agent est un agent chimique.

8. Procédé selon la revendication 1, dans lequel l'agent est une enzyme.

20

9. Procédé selon la revendication 8, dans lequel l'enzyme est une aminopeptidase ou une carboxypeptidase.

10. Procédé selon la revendication 1, dans lequel chaque polypeptide de la pluralité est immobilisé sur un support solide.

25

11. Procédé selon la revendication 10, dans lequel chaque polypeptide est immobilisé dans un réseau.

30

35

40

45

50

55

Figure 1

A

5' d (GAC TTT ACT TGT ACG TGC ATA ATA CGA CTC ACT ATA GGG AGA CTG
AAC ATG GGC AGT CTG AGC CAG ACG CCG GGA CAC AAG GCT GAG CAG CAG
CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAT CAG CAT CAG CAG CAG
CAG CAG CAG CAG CAG CAG CAG CAC CTC ACG AGG GCT CCG GGC CTC ATC
ACC CCG GGT CCC CCC CAC CAG CCC AGC AGA ACC AGT ACG TCC ACA TTT
CAA GTT CAT CAT CAT CAT CAT CAT TGA GAA TCA) -3' (SEQ ID NO. 8)

B

MGSLSQTPGH KAEQQQQQQQ QQQQQHQHQO OOOOOOOOHL
SRAPGLITPG PPGQPSRTST STGQVHHHHH H (SEQ ID NO. 9)

FIGURE 2





FIGURE 3

-Capture of the carboxy terminus.



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 9636732 A [0008] [0069]
- US 5792664 A [0008] [0069] [0230]
- GB 2168478 A [0008]
- US 5777325 A [0047] [0216] [0235]
- US 5742049 A [0047] [0216] [0235]
- US 5654545 A [0047] [0047] [0216] [0216] [0235] [0235]
- US 5641959 A [0047] [0216] [0235]
- US 5760393 A [0047] [0216] [0235] [0235]
- US 649876 A [0066]
- WO 9743617 A [0066]
- US 5643722 A [0092] [0124] [0124] [0165] [0196]
- US 4766072 A [0106]
- US 4983521 A [0117]
- US 5492817 A [0117]
- US 5324637 A [0125]
- US 5604099 A [0129]
- US 5545539 A [0130]
- US 4356270 A [0147]
- US 5547835 A [0147]
- US 5605798 A [0147]
- US 5622824 A [0147]
- WO 9210092 A [0162]
- US 5410068 A [0163]
- US 5612474 A [0163]
- US 5198531 A [0163]
- US 786988 A [0210] [0218]
- US 08787639 B [0210]
- WO 9820166 A [0210] [0218]
- US 08787639 A [0218]
- WO 9324834 A, Chait [0230]
- US 5062935 A [0231]
- WO 9014148 A [0233]
- WO 94121822 A [0238]
- US 5622829 A [0250]
- WO 9213969 A1 [0254]
- EP 0269520 A [0257]
- EP 0655501 A [0257]
- WO 9400562 A [0257]
- US 4683194 A, Mullis [0260]
- WO 19980902 A [0285]
- US 08922201 B [0285]

Non-patent literature cited in the description

- Cooper ; Krawczak. Human Genome Mutations. BI-OS Publ, 1993 [0002]
- Kremer et al. *Science*, 1991, vol. 252, 1711-14 [0003] [0247]
- Fu et al. *Cell*, 1991, vol. 67, 1047-58 [0003] [0247]
- Hirst et al. *J. Med. Genet.*, 1991, vol. 28, 824-29 [0003] [0247]
- Mahadevan et al. *Science*, 1992, vol. 255, 1253-55 [0003] [0247]
- Brook et al. *Cell*, 1992, vol. 68, 799-808 [0003] [0247]
- La Spada et al. *Nature*, 1991, vol. 352, 77-79 [0003]
- Schram. *Mass Spectrometry of Nucleic Acid Components, Biomedical Applications of Mass Spectrometry*, 1990, vol. 34, 203-287 [0008]
- Crain. *Mass Spectrom. Rev.*, 1990, vol. 9, 505-554 [0008]
- Murray. *J. Mass Spectrom. Rev.*, 1996, vol. 31, 1203 [0008]
- Nordhoff et al. *J. Mass Spectrom.*, 1997, vol. 15, 67 [0008]
- Mass Spectrometry. *Meth. Enzymol.* Academic Press, 1990, vol. 193 [0008]
- McLaffery et al. *Acc. Chem. Res.*, 1994, vol. 27, 297-386 [0008]
- Chait ; Kent. *Science*, 1992, vol. 257, 1885-1894 [0008]
- Siuzdak. *Proc. Natl. Acad. Sci., USA*, 1994, vol. 91, 11290-11297 [0008]
- Juhasz et al. *Analysis, Anal. Chem.*, 1996, vol. 68, 941-946 [0047] [0216] [0235]
- Vestal et al. *Rapid Communications in Mass Spectrometry*, 1995, vol. 9, 1044-1050 [0047]
- Hermanson. *Bioconjugate Techniques*. Academic Press, 1996 [0068]
- Huse et al. *Science*, 1989, vol. 246, 1275-1281 [0087]
- Winter ; Harris. *Immunol. Today*, 1993, vol. 14, 243-246 [0087]
- Ward et al. *Nature*, 1989, vol. 341, 544-546 [0087]
- Hilyard et al. *Protein Engineering: A practical approach*. IRL Press, 1992 [0087]
- Borrabeck. *Antibody Engineering*. Oxford University Press, 1995 [0087]
- Harlow ; Lane. *Antibodies: A laboratory manual*. Cold Spring Harbor Laboratory Press, 1988 [0087] [0088] [0089] [0090] [0090] [0095] [0115]

- Sambrook et al. *Molecular Cloning: A laboratory manual*. Cold Spring Harbor Laboratory Press, 1989 [0100] [0104] [0106] [0107] [0114] [0115] [0133]
- Rolff et al. *PCR: Clinical diagnostics and research*. Springer Verlag Publ, 1994 [0100]
- Lin et al. *Nucl. Acids Res.*, 1994, vol. 22, 5220-5234 [0102]
- Jellinek et al. *Biochemistry*, 1995, vol. 34, 11363-11372 [0102] [0103]
- Pagratis et al. *Nature Biotechnol.*, 1997, vol. 15, 68-73 [0102]
- Tam et al. *Nucl. Acids Res.*, 1994, vol. 22, 977-986 [0102]
- Ecker ; Crooke. *BioTechnology*, 1995, vol. 13, 351360 [0102]
- Ausubel et al. *Current Protocols in Molecular Biology*. Green Publ, 1989 [0104]
- Spirin et al. *Science*, 1988, vol. 242, 1162-64 [0111]
- Roberts ; Paterson. *Proc. Natl. Acad. Sci., USA*, 1973, vol. 70, 2330-2334 [0116]
- Anderson. *Meth. Enzymol.*, 1983, vol. 101, 635 [0116]
- Erickson ; Blobel. *Meth. Enzymol.*, 1982, vol. 96, 38 [0117]
- Seong ; RajBhandary. *Proc. Natl. Acad. Sci., USA*, 1987, vol. 84, 334-338 [0121]
- Zubay. *Ann. Rev. Genet.*, 1973, vol. 7, 267 [0125]
- PCR. Newton ; Graham. BIOS Publ. 1994 [0128]
- Wiedmann et al. *PCR Meth. Appl.*, 1994, vol. 3, 57-64 [0128]
- Barany. *Proc. Natl. Acad. Sci., USA*, 1991, vol. 88, 189-93 [0128]
- Walker et al. *Nucl. Acids Res.*, 1994, vol. 22, 2670-77 [0128]
- Higuchi et al. *Bio/Technology*, 1993, vol. 11, 1026-1030 [0128]
- Ellison et al. *J. Biol. Chem.*, 1991, vol. 266, 21150-21157 [0139]
- Narang et al. *Meth. Enzymol.*, 1979, vol. 68, 90 [0147]
- Greene ; Wuts. *Protective Groups in Organic Synthesis*. John Wiley & Sons, 1991 [0160]
- McCray et al. *Ann. Rev. Biophys. Biophys. Chem.*, 1989, vol. 18, 239-270 [0162]
- Rink. *Tetrahedron Letters*, 1976, vol. 28, 3787 [0163]
- Leznoff. *Ace. Chem. Res.*, 1978, vol. 11, 327 [0163]
- Bodansky et al. *Peptide Synthesis*. Academic Press, 1976 [0163]
- Batra et al. *Molecular Immunol.*, 1993, vol. 30, 379-386 [0163]
- Köster et al. *A Versatile Acid-Labile Linker for Modification of Synthetic Biomolecules. Tetrahedron Lett.*, 1990, vol. 31, 7095 [0167]
- Foster. *Organic Charge Transfer Complexes*. Academic Press, 1969 [0168]
- Wentrup. *Reactive Molecules*. John Wiley & Sons, 1984 [0169]
- Fattom et al. *Infect. Immun.*, 1992, vol. 60, 584-589 [0171]
- Welhöner et al. *J. Biol. Chem.*, 1991, vol. 266, 4309-4314 [0171]
- Wong. *Chemistry of Protein Conjugation and Cross-Linking*. CRC Press, 1991 [0174] [0196]
- Hermanson. *Bioconjugate. Techniques*. Academic Press, 1995 [0174]
- Saha et al. *J. Ora. Chem.*, 1993, vol. 58, 7827-7831 [0193]
- Brown et al. *Molec. Divers.*, 1995, 4-12 [0196]
- Rothschild et al. *Nucl. Acids Res.*, 1996, vol. 24, 351-66 [0196]
- Vorm et al. *Anal. Chem.*, 1994, vol. 66, 3281 [0230]
- Valaskovic et al. *Anal. Chem.*, 1995, vol. 67, 3802 [0230]
- Valaskovic et al. *Science*, 1996, vol. 273, 1199-1202 [0232]
- Li et al. *J. Am. Chem. Soc.*, 1996, vol. 118, 1662-1663 [0232]
- Fenn et al. *J. Phys. Chem.*, 1984, vol. 88, 4451-59 [0233]
- Smith et al. *Anal. Chem.*, 1990, vol. 62, 882-89 [0233]
- Ardrey. *Electrospray Mass Spectrometry, Spectroscopy Europe*, 1992, vol. 4, 10-18 [0233]
- Hillenkamp et al. *Matrix Assisted UV-Laser Desorption/Ionization: A New Approach to Mass Spectrometry of Large Biomolecules, Biological Mass Spectrometry*. Elsevier Science Publ, 1990, 49-60 [0233]
- La Spada et al. *Nature*, 1991, vol. 352, 77079 [0247]
- Thompson ; Thompson. *Genetics in Medicine* [0250]
- Nora et al. *Medical Genetics*. Lea and Febiger [0250]
- Nora et al. *Medical Genetics* [0253]
- Edwards et al. *Nucl. Acids Res.*, 1991, vol. 19, 4791 [0254] [0256]
- Beckmann et al. *Genomics*, 1992, vol. 12, 627-631 [0254] [0254]
- Nakamura et al. *Science*, 1987, vol. 235, 1616-1622 [0254]
- Jeffreys et al. *Nature*, 1985, vol. 314, 67-73 [0254]
- Luty et al. *Nucleic Acids Res.*, 1991, vol. 19, 4308 [0254]
- Litt et al. *Nucleic Acids Res.*, 1990, vol. 18, 4301 [0254]
- Litt et al. *Nucleic Acids Res.*, 1990, vol. 18, 5921 [0254]
- Luty et al. *Am. J. Hum. Genet.*, 1990, vol. 46, 776-783 [0254]
- Tautz. *Nucl. Acids Res.*, 1989, vol. 17, 6463-6471 [0254]
- Weber et al. *Am. J. Hum. Genet.*, 1989, vol. 44, 388-396 [0254]
- Polymeropoulos et al. *Nucl. Acids Res.*, 1991, vol. 19, 195 [0256]
- Polymeropoulos et al. *Nucl. Acids Res.*, 1991, vol. 19, 4306 [0256]

- Nishimura et al. *Nucl. Acids Res.*, 1992, vol. 20, 1167 [0256]
- Polymeropoulos et al. *Nucl. Acids Res.*, 1991, vol. 19, 4018 [0256]
- Zuliani et al. *Nucl. Acids Res.*, 1990, vol. 18, 4958 [0256]
- Polymeropoulos et al. *Nucl. Acids Res.*, 1990, vol. 18, 7468 [0256]
- Ploos et al. *Nucl. Acids Res.*, 1990, vol. 18, 4957 [0256]
- Anker et al. *Hum. Mol. Genet.*, 1992, vol. 1, 137 [0256]
- Ratner et al. *Nature*, 1985, vol. 313, 227-284 [0257]
- Wain Hobson et al. *Cell*, 1985, vol. 40, 9-17 [0257]
- Guyader et al. *Nature*, 1987, vol. 328, 662-669 [0257]
- Chakrabarti et al. *Nature*, 1987, vol. 328, 543-547 [0257]
- Gust et al. *Intervirology*, 1983, vol. 20, 1-7 [0257]
- DNA Cloning. 1985, vol. I, II [0260]
- Oligonucleotide Synthesis. 1984 [0260]
- Nucleic Acid Hybridization. 1984 [0260]
- Transcription and Translation. 1984 [0260]
- R.I. Freshney. *Culture of Animal Cells*. Alan R. Liss, Inc, 1987 [0260]
- Immobilized Cells and Enzymes. IRL Press, 1986 [0260]
- B. Perbal. *A Practical Guide to Molecular Cloning*, 1984 [0260]
- Methods In Enzymology. Academic Press, Inc, [0260]
- Gene Transfer Vectors For Mammalian Cells. Cold Spring Harbor Laboratory, 1987 [0260]
- Methods In Enzymology. vol. 154, 155 [0260]
- Immunochemical Methods In Cell And Molecular Biology. Academic Press, 1987 [0260]
- Handbook Of Experimental Immunology. 1986, I-IV [0260]
- Manipulating the Mouse Embryo. Cold Spring Harbor Laboratory press, 1986 [0260]